

# 36th Southern Forest Tree Improvement Conference

---

Overcoming Tree Improvement Bottlenecks  
with New Technologies



**June 7 - 9, 2021**  
Virtual Conference

[36th SFTIC Conference Website](#)

# Table of Contents

---

1. Plenary Session Abstracts.....	pg. 3
2-3. General Session Abstracts	
2A. Pine Genetics and Breeding.....	pg. 7
2B. Forest Health and Restoration.....	pg. 14
3A. Hardwood Genetics and Breeding.....	pg. 21
3B. Forest Health – Pine Pathogens, Insects, and Stress.....	pg. 28
4. Lightning Session Abstracts.....	pg. 34
5. Schatz Symposium Abstracts.....	pg. 52



## Session 1: *Plenary Session*

---

The genetic (and funding) opportunities, challenges and expectations facing tree improvement, even with world-class genomics resources: A *Populus* case study

– *Gerald Tuskan*.....pg. 4

Something in the air – Drone-based high-throughput phenotyping of trees

– *Ingo Ensminger*.....pg. 5

Spectral-based tools for disease resistance phenotyping in trees

– *Anna Conrad*.....pg. 6

# The genetic (and funding) opportunities, challenges and expectations facing tree improvement, even with world-class genomics resources: A *Populus* case study

---

**Gerald A. Tuskan**<sup>1</sup>

<sup>1</sup> Director, The Center for Bioenergy Innovation, Oak Ridge, TN USA 37831 (gtk@ornl.gov)

The current U.S. Administration has a focus on renewable energy, climate change and environmental justice. All of these issues contain forestry related opportunities and challenges. Application of molecular biology and genomics offers potential to ameliorate these problems but the quick delivery window for solutions for these global challenges is confounded by the long reproductive cycles of many long-lived perennial plants and the lack of foundational genomics resources. New sequencing technologies and applications of advanced computing offer hope, but for most species, the realization of that hope is not happening fast enough. In *Populus*, where we have a world-class genomics resources, including high-quality assemblies and annotations for multiple species, fully genotyped association mapping and QTL populations replicated in multiple common gardens, as well as a reliable transformation system based on CRISPR. Using these resources, we have made progress in identifying and validating genes and their associated phenotypes. Examples include EPSP synthase, a novel gene that functions as a transcriptional regulator in the lignin pathway, a lectin kinase that controls colonization by the mycorrhizal fungus *Laccaria bicolor*, and a series of receptor kinases that control Septoria disease resistance. These genes, and many additional SNP markers linked to growth and cell wall traits, are being used in a machine learning approach to stack favorable genes in a genomic selection context. Still, the path to delivering solutions to issues related to renewable energy, climate change and environmental justice remain challenging. Part of the answer is greater funding at all levels, inclusive of federal, state and private sources. Continued improvements in genomics resources, transformation efficiencies and development of early flowering technologies are desperately needed in the conifers and many other hardwood species. The *Populus* case study suggest that rapid progress can be made once such resources are in place. In *Populus*, and all forest tree species, we also need broader field testing of selected and improved plant materials, as well as integrated studies on soil health, carbon sequestration, and water quality. All of us need to become active advocates for the development of these resources. There will be no solution to issues related to renewable energy, climate change and environmental justice without forest trees.

## Something in the air – Drone-based high-throughput phenotyping of trees

---

### ***Ingo Ensminger***

Associate Professor, University of Toronto, Mississauga, ON L5L1C6  
([ingo.ensminger@utoronto.ca](mailto:ingo.ensminger@utoronto.ca))

The field of forest genomics has seen unprecedented advances during the past decade. A suite of genomic resources is now available for enhanced genomic selection. These resources can be used to accelerate breeding cycles and to select genotypes that are better adapted and more resilient to future climate change and diseases. The large-scale phenotyping of populations has become the bottleneck for identifying and connecting the different genomic resources with adaptive traits in populations with thousands of trees. Measuring leaf optical properties using spectral reflectance sensors carried by remotely piloted aircraft systems (drone systems) represent an innovative approach for large-scale phenotyping of tree responses to drought, monitor phenology, and assess differences between tree genotypes in large-scale field experiments. In this presentation I will give an overview of leaf-level, canopy-level and drone based observations of leaf spectral reflectance. I will demonstrate that some of the widely used vegetation indices such as the normalized difference vegetation index (NDVI) and photochemical reflectance index (PRI) vary in their ability to adequately track important traits such as phenology or photosynthetic efficiency. Finally, I will discuss some of the technical challenges of using optical sensors when monitoring complex canopies and why using carotenoid based vegetation indices are particularly useful in order to monitor evergreen conifer canopies.

# Spectral-based tools for disease resistance phenotyping in trees

---

***Anna O. Conrad***<sup>1</sup>

<sup>1</sup> Research Plant Pathologist, USDA Forest Service, Northern Research Station, West Lafayette, IN, USA 47907 (anna.conrad@usda.gov)

Plant specialized metabolites play important roles in tree responses to abiotic and biotic stressors, e.g. plant pathogens and insect pests. For this reason, individual metabolites or groups of metabolites may be useful as biological markers for identifying trees with desired traits, like disease resistance. Infrared spectroscopy is one approach for measuring the complex chemical make-up of tree tissues and has the potential to be used for rapid and high-throughput phenotyping when combined with multivariate statistical analysis or machine learning. Real-time analysis of tree tissues under field conditions is now possible thanks to advances in spectral technology and the availability of low-cost developmental sensor units. Case studies will be presented that demonstrate how infrared spectroscopy can be used to distinguish between disease resistant and susceptible trees. Constraints and opportunities for applying spectral-based tools for tree improvement will also be discussed.



## Concurrent Session 2A: *Pine Genetics and Breeding*

---

- Estimating individual tree heights in loblolly pine biomass plantation using drone imagery and LiDAR-derived measurements  
– *Jessica A. Maynor*.....pg. 8
- Pitro50K: A multispecies genotyping array for tropical pines  
– *Colin Jackson*.....pg. 9
- Application of MateSel software in a loblolly pine breeding program  
– *Trevor D. Walker*.....pg. 10
- Development of elite loblolly pine families for the Piedmont region using clonal progeny testing and genomics  
– *W. Patrick Cumbie*.....pg. 11
- Rangewide patterns of genetic diversity in Table Mountain pine (*Pinus pungens*), an Appalachian endemic  
– *Kevin M. Potter*.....pg. 12
- Variation in salt tolerance of slash pine (*Pinus elliottii*) families  
– *Ronald Schmidting*.....pg. 13

# Estimating individual tree heights in loblolly pine biomass plantation using drone imagery and LiDAR-derived measurements

---

**Jessica A. Maynor**<sup>1</sup>, Steve E. McKeand<sup>2</sup>, Fikret Isik<sup>3</sup>, and James McCarter<sup>4</sup>

<sup>1</sup> Project Leader, ArborGen, Ridgeville, SC, USA 29485 (jamayno@arborgen.com);

<sup>2</sup> Professor of Forestry and Environmental Resources, College of Natural Resources, NC State University, Raleigh, NC 27695.

<sup>3</sup> Professor of Forestry and Environmental Resources, Director of NCSU Advanced Tree Genomics Consortium, and Associate Director of the NC State University Cooperative Tree Improvement Program, College of Natural Resources, NC State University, Raleigh, NC 27695.

<sup>4</sup>Scientific Programmer, Rayonier, Wildlight, FL, USA, 32097.

Height estimation from Unmanned Aerial Systems (UAS) is an ever-increasing interest in natural resource fields. With the use of PhoDAR from drone imagery in combination with LiDAR data, we produced high-resolution digital surface models over two test sites. After individual tree identification, we computed individual tree heights on a young biomass plantation of loblolly pine. PhoDAR derived heights were compared with field measurements. Of the 7,200 study trees, the software was able to detect 7,143 trees. Overall correlations between field measured heights and PhoDAR derived heights among sites was low (0.18). Site 1 had a very low correlation (-0.10), but Site 2 had a very strong correlation (0.79). The difference in correlation was attributed the “bowl effect” commonly noted in some digital elevation models from vertical UAS imagery (James and Robson, 2014). With proper ground control and techniques to mitigate such board-scale deformations, the use of PhoDAR derived measurement can become a more efficient method than traditional field methods.

## Pitro50K: A multispecies genotyping array for tropical pines

---

**Colin Jackson**<sup>1</sup>, Nanette Christie<sup>2</sup>, Melissa Reynolds<sup>2</sup>, Christopher Marais<sup>2</sup>, Madison Caballero<sup>3</sup>, Erik Visser<sup>2</sup>, Sanushka Naidoo<sup>2</sup>, Dominic Kain<sup>4</sup>, Ross W. Whetten<sup>1</sup>, Fikret Isik<sup>1</sup>, Jill Wegrzyn<sup>3</sup>, Gary Hodge<sup>1</sup>, Alexander A. Myburg<sup>2</sup>, Juan Jose Acosta<sup>1</sup>

<sup>1</sup> Department of Forestry & Environmental Resources, North Carolina State University, Raleigh, NC, USA (cajacks7@ncsu.edu; jjacosta@ncsu.edu);

<sup>2</sup> Department of Biochemistry, Genetics and Microbiology, Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, Pretoria, 0002, South Africa;

<sup>3</sup> Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA;

<sup>4</sup>HQPlantations Pty Ltd, North Lakes, LD, 4509, Australia

Using a combination of reduced representation sequencing methods, we performed targeted SNP discovery towards the development of a genome wide, multi-species genotyping array for tropical and subtropical pine species. Pooled RNA-seq data for five species of tropical pines, originating from pathogen challenge experiments, were used to identify transcript based SNP markers. Additionally, target capture sequencing was performed for six species, utilizing pooled DNA from 81 provenances that represented the natural range of the species across Mexico and Central America. Variant calling yielded a total of 1.8 million candidate SNP probesets, 1.3 million and 563K from RNA-seq and target capture respectively. In total, 300K RNA-seq and 120K target capture derived probesets were evaluated on a 420K screening array through the genotyping of 576 trees from eight species, representing the original 81 provenances and commercial breeding material. Evaluation of the screening array probesets resulted in the selection of 50K SNPs for inclusion on the commercial array. These markers included 20K polymorphic SNPs for *P. tecunumanii*, *P. patula*, *P. caribaea*, and *P. oocarpa*, 15K for *P. maximinoi* and *P. greggii*, 13K for *P. elliottii*, and 8K for *P. pseudostrobus*. Of the 50K markers, 75% are polymorphic in two or more species. The Pitro50K genotyping array represents the first high throughput and affordable genomic tool for these species of pines and their hybrids which represent the majority of tropical and subtropical pine plantations globally. Potential applications for the array include population genomics, genetic mapping, molecular breeding, species and hybrid identification, and genetic resource management.

## Application of MateSel software in a loblolly pine breeding program

---

**Trevor D. Walker**<sup>1</sup>, Kitt G. Payn<sup>2</sup>, Steven E. McKeand<sup>2</sup>, and Fikret Isik<sup>2</sup>

<sup>1</sup> NC State University Cooperative Tree Improvement Program, Raleigh, NC, USA 27695-8002  
(trevor\_walker@ncsu.edu);

<sup>2</sup> NC State University Cooperative Tree Improvement Program, Raleigh, NC, USA 27695-8002

The use of the MateSel software is now standard in many livestock and aquaculture breeding programs and is increasingly used in forest tree breeding programs. The software uses a differential evolution algorithm to balance gain and diversity when selecting parents for the next generation and is subject to several logistical constraints that vary based on the biology of the species. The North Carolina State University Cooperative Tree Improvement Program first utilized this algorithm to develop a mating plan for the 4<sup>th</sup>-Cycle breeding program (Isik and McKeand 2019). The Cooperative is now using the software in a novel approach to make forward selections for the 5<sup>th</sup> cycle of breeding by modifying the mating plan output to create a list of progeny test trees to graft for future breeding. We have evaluated several constraints, such as accounting for recent selections in breeding orchards that are not yet producing strobili and accounting for potential selections from families in young progeny tests that are not yet measured. Results indicate that even when the nominal balance of gain and diversity is held constant, considerable impacts on the gain and diversity of the next generation can occur with the inclusion or omission of these logistical breeding constraints.

Isik, F., and S. E. McKeand. 2019. Fourth cycle breeding and testing strategy for *Pinus taeda* in the NC State University Cooperative Tree Improvement Program. *Tree Genetics & Genomes*. 15(5):70.

# Development of elite loblolly pine families for the Piedmont region using clonal progeny testing and genomics

---

**W. Patrick Cumbie**<sup>1</sup>, Jay Coke<sup>1</sup>, Salvador Gezan<sup>2</sup>

<sup>1</sup> ArborGen Inc, Ridgeville, SC, USA 29483 (wpcumbi@arborgen.com);

<sup>2</sup> VSN International Ltd, Hemel Hempstead, United Kingdom

The development timeline to breed, select and commercialize new genetic families for seedling production is a significant bottleneck for forest tree improvement. Reductions in the time required to progeny test and establish seed orchards have been valuable improvements for tree breeding programs in the past. In this study we evaluate the use of clonal progeny testing and genomics to accelerate the development of new selections for the Piedmont region of the southeastern United States. An elite population was created with selections from Alabama, Georgia, North Carolina, and South Carolina. A partial diallel mating design of 22 crosses was completed and 20 to 50 clones per full-sib family were established in 4 locations. After 3 growing seasons, mean height ranged from 10.3 feet to 14.5 feet across the 4 sites with single site  $h^2$  ranging from 0.25 to 0.42. We will present results from current phenotypic and genomic analyses.

## Rangewide patterns of genetic diversity in Table Mountain pine (*Pinus pungens*), an Appalachian endemic

---

**Kevin M. Potter**<sup>1</sup>, Robert M. Jetton<sup>2</sup>, Barbara S. Crane<sup>3</sup>, Valerie D. Hipkins<sup>4</sup>, Craig Echt<sup>5</sup>, W. Andrew Whittier<sup>6</sup>, and Gary Hodge<sup>7</sup>

<sup>1</sup> Research Professor, Department of Forestry and Environmental Resources, North Carolina State University, Research Triangle Park, NC, USA 27719 (kpotter@ncsu.edu);

<sup>2</sup> Associate Professor, Camcore, North Carolina State University, Raleigh, NC, USA 27695;

<sup>3</sup> Regional Geneticist (retired), Southern Region, USDA Forest Service, Atlanta, GA, USA 30309;

<sup>4</sup> Director (former), National Forest Genetics Laboratory, USDA Forest Service, Placerville, CA, USA 95667;

<sup>5</sup> Research Geneticist (retired), Southern Institute of Forest Genetics, USDA Forest Service, Saucier, MS, USA 39574;

<sup>6</sup> Research Associate, Camcore, North Carolina State University, Raleigh, NC, USA 27695;

<sup>7</sup> Professor, Camcore, North Carolina State University, Raleigh, NC, USA 27695

Table Mountain pine (*Pinus pungens*) is an imperiled tree species endemic to the Appalachian Mountains from Georgia to Pennsylvania. Generally reliant on fire for regeneration, its formerly fragmented but widespread distribution has dwindled to fewer than 30,000 acres. It typically occurs in small and geographically isolated populations at elevations of 1,000-4,000 feet in rocky soils on south- and west-facing ridgelines. The suppression of wildfire has allowed hardwoods to take over many sites formerly dominated by Table Mountain pine. We quantified the genetic diversity of 346 trees in 32 populations across the range of the species using data from seven highly polymorphic simple sequence repeat (SSR) loci. The species is relatively inbred ( $F_{IS} = 0.208$ ) while differentiation among populations was relatively low ( $F_{ST} = 0.029$ ). Areas with high genetic diversity included the Blue Ridge Mountains of west-central Virginia and of southwest Virginia and northwestern North Carolina, and the Great Smoky Mountains along the Tennessee-North Carolina border. Populations with unique alleles were scattered throughout the species range. We found a strong negative correlation between population isolation and heterozygosity and a strong positive correlation between population isolation and genetic differentiation. Few significant differences in genetic diversity metrics existed among the seed collection zones proposed for Table Mountain pine, except that populations in the most northern seed zone had lower heterozygosity than the southern seed zones, and that central seed zone populations are less genetically differentiated than the northern and southern seed zones. Genetic structure analysis suggested the existence of six to eight genetic clusters in Table Mountain pine. Some of these genetic clusters were associated with different parts of the species distribution. These results add to our limited knowledge of genetic variation across the distribution of this rare and threatened conifer.

## Variation in salt tolerance of slash pine (*Pinus elliottii*) families

---

**R.C. Schmidting**<sup>1</sup>, James Roberds<sup>1</sup>, and C. Dana Nelson<sup>2</sup>

<sup>1</sup>Scientist Emeritus, <sup>2</sup> Project Leader, Southern Institute of Forest Genetics, USDA Forest Service, Saucier, MS, USA 39574 (rschmidting@gmail.com)

The average temperature is going up, the glaciers are melting, and sea levels are rising! Salt tolerance will become an important factor in restoring vegetation in coastal environments. On the Gulf Coast of Mississippi, Alabama, and Florida, slash pine (*Pinus elliottii*) occurs in coastal and barrier island environments that are often subject to salt-water exposure. We tested 1-0 seedlings of open-pollinated families of slash pine from three populations: a barrier island (20 km offshore), a near-shore island (100 m off-shore) and inland (55 km north of the coast) populations. After trying many treatment combinations and concentrations of salt-water we found that three treatments of watering from below at bi-weekly intervals with natural brackish water (11 to 20 ppt salinity) from the Mississippi Sound gave us overall survival around 50% and good family separation. We did not find significant ecotypic variation among seed sources, but we did find ample and significant family-in-source variation in survival. Survival of families varied from 0 to 70%, a range that appears to be useable in a breeding program.



## Concurrent Session 2B: *Forest Health and Restoration*

---

- Conservation and population genetics of the federally endangered *Torreya taxifolia* (Taxaceae)  
– *Lauren Eserman*.....pg. 15
- Evolutionary origins of the hemlock woolly adelgid resistance-disparity between Asian and North American hemlocks (*Tsuga* sp.).  
– *Karl C. Fetter*.....pg. 16
- Development of a new genetic transformation system for white and green ash using embryogenic cultures  
– *A. Ryan Tull*.....pg. 17
- Mapping QTLs for blight resistance and morphological traits with complex chestnut families  
– *Shenghua Fan*.....pg. 18
- Evidence of Fraser fir partial resistance to the balsam woolly adelgid  
– *Austin M. Thomas*.....pg. 19
- Saving butternut  
– *Jeanne Romero-Severson*.....pg. 20

## Conservation and population genetics of the federally endangered *Torreya taxifolia* (Taxaceae)

---

**Lauren A. Eserman**<sup>1</sup>, Ashlynn Smith<sup>2</sup>, Laurie Blackmore<sup>3</sup>, Juliet Rynear<sup>4</sup>, Lilly Anderson-Messecc<sup>5</sup>, Jim Leebens-Mack<sup>6</sup>, Chazz Jordan<sup>7</sup>, Emily E. D. Coffey<sup>8</sup>

<sup>1</sup> Research Scientist, Conservation & Research Department, Atlanta Botanical Garden, Atlanta, GA, USA 30309 (leserman@atlantabg.org);

<sup>2</sup> Gulf Coast Coordinator, Research Scientist, Conservation & Research Department, Atlanta Botanical Garden, Atlanta, GA, USA 30309;

<sup>3</sup> Conservation & Research Manager, Conservation & Research Department, Atlanta Botanical Garden, Atlanta, GA, USA 30309;

<sup>4</sup> Executive Director, Florida Native Plant Society, Melbourne, FL, USA 32902-0278;

<sup>5</sup> TorreyaKeepers Project Director, Florida Native Plant Society, Melbourne, FL, USA 32902;

<sup>6</sup> Professor and Interim Curator of the University of Georgia Herbarium, Plant Biology Department, University of Georgia, Athens, GA, USA 30602;

<sup>7</sup> Graduate Student, Dept of Plant Biology, University of Georgia, Athens, GA, USA 30602;

<sup>8</sup> Vice President for Conservation and Research, Conservation & Research Department, Atlanta Botanical Garden, Atlanta, GA, USA 30309.

Florida torreya, or *Torreya taxifolia* (Taxaceae), is one of the most endangered conifers in North America and is endemic to the ravines east of the Apalachicola river in the Florida panhandle. In the last century, populations have declined from nearly 700,000 trees in the early 1900s to around 700 trees today. This dramatic decline is the result of an invasive fungal pathogen, *Fusarium torreyae*. For the last 30 years, staff at the Atlanta Botanical Garden have been collecting cuttings for safeguarding, caging trees to prevent deer browsing, and tagging and monitoring wild trees in Torreya State Park and the Nature Conservancy's Apalachicola Bluffs and Ravines Preserve. Recently, partnership with the Florida Native Plant Society has allowed outreach to private landowners to locate, tag, and collect cuttings of trees on private lands. Using the collection of trees from across its range, we are performing conservation genetic studies of *Torreya taxifolia* using target gene capture to assess the level of genetic diversity and population structure remaining in the wild. Conifers are notorious for having extremely large and highly repetitive nuclear genomes, making typical population genomic techniques such as Genotype-by-Sequencing (GBS), restriction site-associated DNA sequencing (RADseq), and genome skimming unfeasible. Gene capture, in contrast, allows for targeted sequencing of specific loci in the genome and allows us to overcome the problem of the large, repetitive genome structure. This method is a cost-effective way to obtain DNA sequence variation necessary to distinguish among closely related populations. Together, these projects will advance conservation efforts for this critically imperiled conifer.

# Evolutionary origins of the hemlock woolly adelgid resistance-disparity between Asian and North American hemlocks (*Tsuga* sp.).

---

**Karl C. Fetter**<sup>1</sup>, John R. Butnor<sup>2</sup>, Katherine E. Smtih<sup>3</sup>, Albert G. Abbott<sup>4</sup>, and C. Dana Nelson<sup>5</sup>

<sup>1</sup> Postdoctoral Research Associate, University of Georgia, Athens, GA, USA 30602 (kf@uga.edu)

<sup>2</sup> Research Plant Physiologist, USDA Forest Service, Southern Research Station, Burlington, VT, 05405;

<sup>3</sup> Biological Science Technician, Southern Research Station, USDA Forest Service, Gainesville, FL, USA 32611;

<sup>4</sup> Bioscience Team Leader, Forest Health Research and Education Center, University of Kentucky, Lexington, KY, USA 40506;

<sup>5</sup> Project Leader, USDA Forest Service, Southern Research Station, Forest Health Research and Education Center, Lexington, KY, USA 40506 (charles.d.nelson@usda.gov)

Evolutionary theory provides a framework for understanding population and species level variation, which, when translated into a context for understanding the scope and magnitude of a conservation problem, can provide novel solutions. The hemlock woolly adelgid (HWA, *Adelges tsugae*) is a sap-feeding pest evolved in eastern Asia and introduced into Eastern North America in the 1950's that has nearly extirpated local populations of the eastern hemlock (*Tsuga canadensis*) across the central and southern portions of its range. The Chinese hemlock (*T. chinensis*), in contrast, has a long evolutionary history of exposure to the HWA and is largely resistant. Here, we seek to use evolutionary theory to understand the origins of the disparity of resistance to HWA by focusing on patterns of allele frequency variation in terpenoid biosynthesis pathway genes. Terpenoids are generally thought to be stress avoidance or tolerance mechanisms in gymnosperms, and are likely candidates to explain patterns of resistance to the HWA within *Tsuga*. We employ neutrality tests in gene coding regions to understand patterns of sequence evolution. This study will provide an important context for understanding the origin of epidemics after the introduction of novel pests as well as provide important genomic resources for an understudied and imperiled tree of North American forests.

# Development of a new genetic transformation system for white and green ash using embryogenic cultures

---

**A. Ryan Tull**<sup>1</sup>, Heather J. Gladfelter<sup>2</sup>, Flavia Pampolini<sup>3</sup>, Shenghua Fan<sup>4</sup>, Lynne Rieske-Kinney<sup>5</sup>, C. Dana Nelson<sup>6</sup>, Albert G. Abbott<sup>7</sup>, and Scott A. Merkle<sup>8</sup>

<sup>1</sup> Research Professional, University of Georgia, Athens, GA, USA 30602 (rtull@uga.edu);

<sup>2</sup> Postdoctoral Associate, University of Georgia, Athens, GA, USA 30602;

<sup>3</sup> Graduate Research Assistant, University of Kentucky, Lexington, KY USA 40506;

<sup>4</sup> Postdoctoral Associate, University of Kentucky, Lexington, KY USA 40506;

<sup>5</sup> Professor, University of Kentucky, Lexington, KY USA 40506;

<sup>6</sup> Project Leader, USDA Forest Service, University of Kentucky, Lexington, KY USA 40506;

<sup>7</sup> Adjunct Faculty, University of Kentucky, Lexington, KY USA 40506;

<sup>8</sup> Professor, University of Georgia, Athens, GA, USA 30602

All North American ash species are under threat from the emerald ash borer (EAB; *Agrilus planipennis*), an exotic wood-boring beetle that has already destroyed millions of ash trees in the U.S. and Canada. One potential approach to managing this destructive insect is to develop a host-induced gene silencing strategy employing transgenic ash that express RNA interference (RNAi) constructs targeting EAB-specific genes. Exogenous double-stranded RNA triggers the RNAi pathway and can silence key genes, disrupting protein function and causing insect mortality. An important prerequisite for advancing this technology is a reliable transformation/regeneration system for ash. We tested embryogenic cultures of white ash and green ash as target material for *Agrobacterium*-mediated gene transfer with marker genes. In a series of experiments, cells of different suspension-grown white ash and green ash embryogenic cultures were collected and inoculated with *A. tumefaciens* strain AGL-1 carrying the pFHI-GUSi expression vector, which, in addition to the intron-GUS reporter gene, also includes the nptII selectable marker gene. After inoculation, cells were maintained on semi-solid medium that contained 35 mg/l geneticin and 300 mg/l timentin to select for transformed cells and eradicate *Agrobacterium*. Once putative transgenic events were identified, they were assayed at every developmental stage from callus to plantlets using the fluorometric assay for  $\beta$ -glucuronidase to confirm the presence of the transgene. When plants were obtained, DNA was isolated from leaf tissue and PCR was used to confirm the insertion of the intron-GUSi gene. To date, two genotypes each of white ash and green ash have been transformed and multiple events from both species have been successfully acclimated to soil.

# Mapping QTLs for blight resistance and morphological traits with complex chestnut families

---

**Shenghua Fan**<sup>1,2</sup>, Laura L. Georgi<sup>3</sup>, Frederick V. Hebard<sup>4</sup>, Jiali Yu<sup>5</sup>, Albert G. Abbott<sup>6</sup>, C. Dana Nelson<sup>7,8</sup>

<sup>1</sup> Research associate, Forest Health Research and Education Center, University of Kentucky, Lexington, KY, USA

<sup>2</sup> Research associate, Department of Horticulture, University of Kentucky, Lexington, KY 40546

<sup>3</sup> Research scientist, Forest Health Research and Education Center, Damascus, VA, USA

<sup>4</sup> Chief Scientist Emeritus, The American Chestnut Foundation, Meadowview, VA, USA

<sup>5</sup> Graduate research assistant, UT-ORNL Graduate School of Genome Science and Technology, Knoxville, TN 37996

<sup>6</sup> Bioscience team leader, Forest Health Research and Education Center, University of Kentucky, Lexington, KY, USA

<sup>7</sup> Co-Director, Forest Health Research and Education Center, USDA Forest Service, Southern Research Station, Lexington, KY, USA

<sup>8</sup> Project leader, Southern Institute of Forest Genetics, USDA Forest Service, Southern Research Station, Saucier, MS, USA

Chestnut blight (caused by *Cryphonectria parasitica*, *Cp*), together with *Phytophthora* root rot, has nearly extirpated American chestnut (*Castanea dentata*). In contrast to the susceptibility of American chestnut, many Chinese chestnut (*C. mollissima*) genotypes are resistant to blight. In this research, we performed a series of GWAS studies for blight resistance originating from three unrelated Chinese chestnut trees (Mahogany, Nanking and M16) and a QTL study on a Mahogany-derived F2 family. We evaluated trees for resistance to blight after artificial inoculation and scored nine morphological traits that are the hallmarks of species differentiation between American and Chinese chestnuts. Blight resistance and morphological trait QTLs that were identified in the GWAS and F2 mapping studies were compared. Results support a moderately complex genetic architecture for blight resistance, as 13 QTLs were found on 11 chromosomes across all studies, including two previously identified QTLs (Cbr1 and Cbr3). Additionally, blight resistance QTLs overlapped with 9 of 15 morphological trait loci indicating that it will be challenging, but still possible, to eliminate Chinese chestnut alleles for these distinguishing traits while still achieving high blight resistance in the backcross hybrid chestnuts. Finally, comparison between QTL intervals for blight resistance and those previously published for root rot resistance, revealed common disease resistance loci on chromosome A, E, and K.

## Evidence of Fraser fir partial resistance to the balsam wooly adelgid

---

**Austin M. Thomas**<sup>1</sup>, Benjamin C. Smith<sup>2</sup>, Ross Whetten<sup>3</sup>, Jill L. Wyzern<sup>4</sup>, Lilian P. Matallana-Ramirez<sup>5</sup>, John Frampton<sup>6</sup>, David C. Tillotta<sup>3</sup>, Robert M. Jetton<sup>7</sup>

<sup>1</sup> Ph.D. Candidate, North Carolina State University, Raleigh, NC, USA 27695  
(amthom22@ncsu.edu)

<sup>2</sup> Research Scholar, North Carolina State University, Raleigh, NC, USA 27695

<sup>3</sup> Professor, North Carolina State University, Raleigh, NC, USA 27695

<sup>4</sup> Professor, University of Connecticut, Storrs, CT, USA 06269

<sup>5</sup> Research Associate, North Carolina State University, Raleigh, NC, USA 27695

<sup>6</sup> Professor Emeritus, North Carolina State University, Raleigh, NC, USA 27695

<sup>7</sup> Associate Professor, North Carolina State University, Raleigh, NC, USA 27695

The balsam woolly adelgid, *Adelges piceae* (BWA), was first reported on Fraser fir, *Abies fraseri*, on Mount Mitchell in 1955. A novel herbivore in North America, BWA was responsible for sharp declines in wild Fraser fir populations in the 1960's and 70's. Tree mortality is thought to be the result of systemic rotholz formation. Rotholz is dense and resinous wood that restricts water transport throughout the tree. Fraser fir is the premier Christmas tree species in North Carolina and plantations require frequent pesticide applications to prevent BWA related damage and mortality. Breeding Fraser fir for BWA tolerance or resistance traits may lessen or eliminate the need for pesticide treatments in the future. In this genetic study we evaluated 37 clones of improved Fraser fir for increased tolerance or resistance to BWA using tree growth, gouting, loss of apical dominance, systemic rotholz formation, and mortality as performance metrics. Additionally, we evaluated the lateral bud transcriptome and foliar terpenoid expression of select clones in response to BWA feeding. We found a broad range of responses among clones ranging from extreme susceptibility to significant levels of resistance. Increased foliar concentration of several terpenoid compounds in response to BWA feeding were observed across all clones but this response did not correlate with clone performance. Significant upregulation of genes related to plant defense, lignin synthesis, and traumatic resin duct formation observed in high performing clones, however, does suggest a targeted defensive response to BWA feeding. We believe these data provide strong evidence that some degree of BWA resistance is present in Fraser fir populations which could be further enhanced through selective breeding.

## Saving butternut

---

Carolyn C. Pike<sup>1</sup>, Martin Williams<sup>2</sup>, Andrea Brennan<sup>3</sup>, Keith Woeste<sup>4</sup>, James Jacobs<sup>5</sup>, Sean Hoban<sup>3</sup>, Melanie Moore<sup>5</sup>, **Jeanne Romero-Severson**<sup>6</sup>

<sup>1</sup> USDA Forest Service Eastern Region State and Private Forestry, West Lafayette IN

<sup>2</sup> Canadian Forest Service, Ottawa ON

<sup>3</sup> The Morton Arboretum, Lisle IL

<sup>4</sup> USDA Forest Service Northern Research Station, West Lafayette IN

<sup>5</sup> USDA Forest Service Forest Health Protection, St Paul MN

<sup>6</sup> Professor, University of Notre Dame, South Bend, IN 46556

Butternut is a relatively uncommon hardwood tree native to eastern North America. Butternut abundance has declined over the past fifty years, primarily due to the invasive pathogen *Ophiognomonia clavigignenti-juglandacearum* and loss of suitable habitat for regeneration. Despite steep population declines, genetic diversity range-wide remains fairly high. While there is little evidence for even moderate resistance in native butternut, hybrids with Japanese walnut, a closely related species, display enough resistance to persist on the landscape without protection and bear abundant nut crops year after year. Cryostorage of native embryogenic axes has yielded promising initial results as a strategy for gene conservation, but additional action is needed to conserve the remaining native gene pool. We describe a practical recurrent selection strategy for resistance breeding in butternut, using sources of resistance from naturally occurring hybrids, hybrids in research orchards, sources of native trees from as many regions as possible, and targeted genotyping.



## Concurrent Session 3A: *Hardwood Genetics and Breeding*

---

- White oak genetics and tree improvement program: Range-wide collaborative effort, early results, and lessons learned  
– *Laura DeWald*.....pg. 22
- Assembly and annotation of the green ash genome  
– *Matthew Huff*.....pg. 23
- Fast-track transformation and flowering for rapid-cycle breeding in poplar  
– *C.J. Tsai*.....pg. 24
- Exploring the genetic diversity of an extinct tree species in the wild, *Franklinia alatamaha*, using genotyping-by-sequencing of the surviving cultivated population  
– *Heather J. Gladfelter*.....pg. 25
- CRISPR/Cas9-knockout of trichome-regulating MYBs in *Populus* alter light sensitivity and wax composition  
– *William P. Bewg*.....pg. 26
- Ribosomal DNA organization and genomic composition of satellite chromosome in American (*Castanea dentata* (Marshall) bork.) And Chinese (c. *Mollissima blume*) chestnut  
– *Nurul Islam-Faridi*.....pg. 27

## White oak genetics and tree improvement program: Range-wide collaborative effort, early results, and lessons learned

---

**Laura E. DeWald**<sup>1</sup>, Zachary J. Hackworth<sup>2</sup>, and C. Dana Nelson<sup>3</sup>

<sup>1</sup> Tree Improvement Specialist and Adjunct Professor, University of Kentucky, Lexington, KY USA 40546 (Laura.DeWald@uky.edu);

<sup>2</sup> Research Analyst, University of Kentucky, Lexington, KY, USA 40546;

<sup>3</sup> Research Geneticist and Project Leader, USDA Forest Service, Southern Research Station, Lexington, KY USA 40546 and Southern Institute of Forest Genetics, Saucier, MS USA 39574

The role of genetics and tree improvement is a focus area of the White Oak Initiative in recognition of the importance this work plays in our ability to respond to increasing pressures on the white oak resource. The White Oak Genetics and Tree Improvement program (WOGTIP) ([www.white-oak-genetics.ca.uky.edu](http://www.white-oak-genetics.ca.uky.edu)) is collaborating with industry and forestry, conservation, and wildlife agencies and organizations to: quantify white oak genetic variation and improve traits that have economic and ecological value. WOGTIP is working toward providing a sustainable supply of improved white oak seedlings to meet current and future demands that will improve our ability to conserve and restore white oak in the forest to achieve a variety of ecological, conservation and economic goals at regional and national levels. WOGTIP is also providing genetic resources for academic and industrial research and development. The plan is organized as three phases: Collecting Genetic Material, Progeny Testing, and Seed Production /Orchard Establishment. WOGTIP is coordinated through University of Kentucky, but is being implemented through a range-wide network of volunteers, partnerships and collaborations. Volunteers, and state and federal agency personnel have collected acorns and scions, seedlings are grown at a Kentucky Division of Forestry state nursery, and progeny tests, seed production areas, and grafted seed orchards are (or will be) established on private and public lands throughout the eastern US. Acorn collections representing half-sibling families of 190 parent trees have been made over the past two years. Seedlings from 63 of the acorn collections made in 2019 were outplanted into the first progeny tests in April 2021. Early results show genetic variation among and within these half-sibling families in acorn size, seedling height and apical dominance, root biomass and morphology, and spring phenology.

## Assembly and annotation of the green ash genome

---

**Matthew Huff**<sup>1\*</sup>, Josiah Seaman<sup>2,3</sup>, Di Wu<sup>4</sup>, Tetyana Zhebentyayeva<sup>4</sup>, Laura Kelly<sup>2,3</sup>, Nurul Faridi<sup>5,6</sup>, C. Dana Nelson<sup>6,7</sup>, Endymion Cooper<sup>2</sup>, Enrico Bonello<sup>8</sup>, Jennifer Koch<sup>9</sup>, Jeanne Romero-Severson<sup>10</sup>, John E. Carlson<sup>4</sup>, Richard Buggs<sup>2,3</sup> and Margaret Staton<sup>1</sup>

<sup>1</sup> Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, Tennessee, USA

<sup>2</sup> School of Biological and Chemical Sciences, Queen Mary University of London, London, UK

<sup>3</sup> Royal Botanic Gardens, Kew, Richmond, Surrey, UK

<sup>4</sup> Department of Ecosystem Science and Management, Pennsylvania State University, University Park, Pennsylvania, USA.

<sup>5</sup> Department of Ecosystem Science and Management, Texas A&M University, College Station, Texas, USA.

<sup>6</sup> USDA Forest Service, Southern Research Station, Saucier, Mississippi, USA

<sup>7</sup> Forest Health Research and Education Center, University of Kentucky, Lexington, KY, USA

<sup>8</sup> Department of Plant Pathology, The Ohio State University, Columbus, Ohio, USA

<sup>9</sup> United States Department of Agriculture, Forest Service, Northern Research Station, Delaware, Ohio, USA

<sup>10</sup> Department of Biological Sciences, Notre Dame University, 46556 Notre Dame, Indiana, USA

Once the most widely distributed ash tree in North America, the green ash (*Fraxinus pennsylvanica*) has faced high mortality as a result of the non-native invasive emerald ash borer (EAB; *Agrilus planipennis*). A small percentage of native green ash trees that remain healthy in long-infested areas, termed “lingering ash,” display partial resistance to the insect, indicating that breeding and propagating populations with higher resistance to the pest may be possible. Genomic resources could assist green ash resistance breeding programs by enabling identification of genetic markers associated with resistance and other important traits. However, at present only scaffold-level genome assemblies, without gene or trait information, are available. As a first step toward providing information needed by tree breeders, we developed a 757 Mbp chromosome-scale assembly scaffolded with a newly expanded genetic linkage map containing over 4,000 SNP markers. Gene annotation of the assembled genome sequence identified 35,470 protein coding genes. Synteny and base-pair substitution analysis confirmed the presence of the previously reported Oleaceae family whole genome duplication. Interestingly, the chromosomes are only moderately rearranged since the duplication event and residual syntenic blocks are identified both within the green ash genome and between green ash and olive. In addition, we demonstrate further utility of the new assembly through reference-guided scaffolding of publicly available genomes from 28 other *Fraxinus* species and subspecies. These resources and analyses provide a new opportunity to characterize EAB response among species spanning from resistant to susceptible in phenotype and should benefit EAB-resistance breeding programs and ash restoration efforts.

## Fast-track transformation and flowering for rapid-cycle breeding in poplar

---

Ran Zhou, Bindu Simon, Margot S.S. Chen, Maria A. Ortega, W. Patrick Bewg and  
***Chung-Jui Tsai***

Warnell School of Forestry and Natural Resources, Department of Plant Biology and  
Department of Genetics, University of Georgia, Athens, GA 30606, USA (cjtsai@uga.edu)

Forest trees have a juvenile phase of several years to several decades. The long generation time is a major impediment to breeding and a bottleneck to fundamental research involving reproductive traits, transgenerational monitoring, and genetic containment. Several regulators of floral development, such as *LEAFY* and *FLOWERING LOCUS T* (*FT*), have been used to induce precocious flowering in annual models. Translating these findings into the poplar system has met with various challenges, including dwarfism and sterile flowers. While the use of heat-inducible promoters for floral gene expression has alleviated many of the developmental anomalies, tedious heat treatments over multiple weeks and months are detrimental to microsporogenesis. The efficacy is also season- and genotype-dependent, limiting widespread adoption of this method. For dioecious species like *Populus*, early flowering male and female genotypes must be available for cross-pollination. Building on the recent discovery of a single female-specific cytokinin response regulator (*FRR*) underlying sex determination in *Populus*, we reasoned that early flowering and sex switch can be engineered simultaneously to produce male, female and/or hermaphrodite individuals from the same genetic background to accelerate poplar breeding and research. We used CRISPR/Cas9 to knock out a negative regulator of floral initiation in a female hybrid aspen *Populus tremula* x *alba* INRA 717-1B4. *In vitro* flowering was readily observed within 3-4 months of *Agrobacterium*-mediated transformation. When *FRR* was also targeted, transgenic plants developed male and sometimes hermaphrodite flowers in tissue culture. With further development of an efficient early flowering and sex switch system like this, the promise of CRISPR to accelerate rapid-cycle breeding or rapid-cycle genomic selection via controlled crosses is finally within reach for long-lived woody perennials.

Exploring the genetic diversity of an extinct tree species in the wild,  
*Franklinia alatamaha*, using genotyping-by-sequencing of the surviving  
cultivated population

---

**Heather J. Gladfelter**<sup>1</sup>, Lav Yadav<sup>2</sup>, Scott A. Merkle<sup>3</sup>, and H. Dayton Wilde<sup>4</sup>

<sup>1</sup> Postdoctoral Associate, University of Georgia, Athens, GA, USA 30602  
(hjgladf@uga.edu);

<sup>2</sup> Graduate Research Assistant, University of Georgia, Athens, GA USA 30602

<sup>3</sup> Professor, University of Georgia, Athens, GA, USA 30602

<sup>4</sup> Professor, University of Georgia, Athens, GA, USA 30602

Genetic diversity of an extinct tree species in the tea family (Theaceae), *Franklinia alatamaha*, has been of curious interest since this tree only existed as one remote population when growing in its native habitat on the Georgia Coastal Plain. *F. alatamaha* was first documented by American naturalists John and William Bartram during their travels to Florida in 1765. The only known population of *F. alatamaha* became extinct in the wild by the early 1800s. Fortunately, William Bartram collected seed from this population in 1773 and brought it back to Philadelphia, where he grew plants from the harvested seed, which has allowed *F. alatamaha* to be cultivated as an ornamental for nearly 250 years. All extant *F. alatamaha* trees in the world are derived from that seed collection. The only opportunity to infer the genetic variation that once existed in the original population was to examine the genetic variation in the cultivated population. Leaves from live and herbarium accessions were obtained from 42 sites worldwide. Genotyping-by-sequencing (GBS) was used to determine the genetic diversity and structure of 76 *F. alatamaha* accessions, including a 178-year-old herbarium specimen. STRUCTURE analysis with 9604 high-quality single-nucleotide polymorphisms (SNPs) identified two subpopulations within the cultivated accessions. This result was supported by UPGMA (unweighted pair group method with arithmetic mean) and principal component analyses. F statistics indicated that there was a moderate level of genetic diversity among the cultivated accessions ( $F_{ST} = 0.09$ ), with more genetic diversity among accessions within a subpopulation than between the two subpopulations. The inbreeding coefficient of the cultivated accessions was low ( $F_{IS} = -0.4902$ ), indicating that the sampled trees represent what was once a highly outcrossing population. The genetic differentiation identified in this study may be useful for further development of new horticultural traits such as disease resistance to *Phytophthora* root rot, which inhibits *F. alatamaha* from being cultivated in the region to which it was once native.

# CRISPR/Cas9-knockout of trichome-regulating MYBs in *Populus* alter light sensitivity and wax composition

---

**William P. Bewg**<sup>1</sup>, Margot Chen<sup>1</sup>, Scott Harding<sup>1</sup>, Jingyin Yu<sup>1</sup>, Ran Zhou<sup>1</sup>, Timothy Tschaplinski<sup>2</sup>, Nancy Engle<sup>2</sup>, Braz Vaidya<sup>3</sup>, Nirmal Joshee<sup>3</sup> and Chung-Jui Tsai<sup>1</sup>

<sup>1</sup> School of Forestry and Natural Resources, Department of Genetics, and Department of Plant Biology, University of Georgia, Athens, Georgia, USA 30602 (bewg@uga.edu);

<sup>2</sup>Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA 37830;

<sup>3</sup>Fort Valley State University, Fort Valley, Georgia, USA 31030.

In all habitats, plants face a variety of environmental stresses, and thus, have evolved an array of physiological and transcriptional coping mechanisms. Acting as a physical barrier between plant and its environment, hair-like trichomes act multifunctionally to provide pest defense and UV shielding whilst reducing transpiration rates. Additionally, they can also act as a locale for secondary metabolite synthesis and storage. The role and regulation of trichomes is well studied in herbaceous models, but less so in poplar, a woody perennial with bioenergy importance. Previous research identified the transcription factor *PtaMYB186* as a positive regulator of trichome initiation during early stages of leaf development in *Populus tremula* x *P. alba* (IRNA 717-1B4). Here, the CRISPR/Cas9 system was utilized to target *PtaMYB186* and its close paralogs for knockout mutagenesis in poplar. The regeneration of trichomeless mutants confirmed the regulatory roles of the MYB transcription factors during trichome initiation. These trichomeless poplar had increased pest susceptibility, though unexpectedly, growth and leaf transpiration rates were not affected. Additionally, light-regulated genes were found to be differentially expressed and exposing the trichomeless mutants to a high-light environment significantly increased synthesis of anthocyanins, a class of known photoprotective metabolites. Notably, cuticle wax and whole leaf analyses found a complete absence of triterpenes in the mutants, suggesting biosynthesis and storage of triterpenes in poplar occurs in the trichomes. Together, these findings contribute further insights into the multifunctional role of trichomes in poplar as both a pest and light barrier, as well as a site of triterpene biosynthesis and storage.

# Ribosomal DNA organization and genomic composition of satellite chromosome in American (*Castanea dentata* (Marshall) bork.) and Chinese (*c. Mollissima* blume) chestnut

---

**Nurul Islam-Faridi**<sup>1</sup>, Tetyana Zhebentyayeva<sup>2</sup>, George L Hodnett<sup>3</sup>, Laura L Georgi<sup>4</sup>, Paul H Sisco<sup>5</sup>, Frederick V Hebard<sup>4</sup>, and C Dana Nelson<sup>6,7</sup>

<sup>1</sup> USDA Forest Service, Forest Tree Molecular Cytogenetics Laboratory, Southern Institute of Forest Genetics, Southern Research Station, College Station, TX 77843 USA

<sup>2</sup> The Schatz Center for Tree Molecular Genetics, Department of Ecosystem Science and Management, The Pennsylvania State University, University Park, PA 16802 USA

<sup>3</sup> Dept. of Soil & Crop Sciences, Texas A&M University, College Station, TX 77843 USA

<sup>4</sup> Meadowview Research Farms, The American Chestnut Foundation, 29010 Hawthorne Drive, Meadowview, VA 24361 USA

<sup>5</sup> The American Chestnut Foundation, 50 North Merrimon Ave., Suite 115, Asheville, NC 28804 USA

<sup>6</sup> Forest Health Research and Education Center, USDA Forest Service, Southern Research Station, Lexington, KY 40546 USA

<sup>7</sup> USDA Forest Service, Southern Institute of Forest Genetics, Harrison Experimental Forest, 23332 Success Road, Saucier, MS 39574 USA

The American chestnut once known as the "King of the Appalachian Forest" has been decimated by a dreadful disease, chestnut blight, caused by a non-native invasive fungus, *Cryphonectteria parasitica*. Chinese chestnut (CC), in contrast, is resistant to blight and efforts are being made to transfer this trait into American chestnut (AC) through backcross breeding and genetic engineering. To facilitate these efforts CC has been genetically mapped and sequenced; however, cytogenetic information is generally lacking in these species. Fluorescence in situ hybridization (FISH) is a modern cytogenetic technique capable of revealing the structural organization of genomes by physically locating molecular markers and/or genes on specific chromosome(s). Using FISH we have identified two 35S rDNA (one major and one minor), and one 5S rDNA loci in AC and CC. A second 5SrDNA locus has also been identified in CC, which is co-localized distally to the major 35S rDNA locus. The major 35S rDNA loci of AC and CC are located in terminal and sub-terminal positions, respectively; and their associated satellites are structurally and organizationally different from each other. The AC satellite is very small and always obscured by the 35S rDNA signal, whereas the CC satellite constitutes about 25% of the total length of the chromosome and only the proximal half is obscured by the 35S signal. The chromosomal differences reveal evolutionary divergence between AC and CCs and highlight potential genetic challenges when using interspecific populations for breeding purposes.



## Concurrent Session 3B: *Forest Health – Pine Pathogens, Insects, and Stress*

---

- Rapid-phenotyping of fusiform rust disease resistance in loblolly pine through vibrational spectroscopy  
– *Simone Lim-Hing*.....pg. 29
- NLR diversity and candidate fusiform rust resistance genes in loblolly pine  
– *Daniel Ence*.....pg. 30
- Longitudinal study on in-field trial rust resistance and seedling screening results  
– *Stephanie Fong*.....pg. 31
- Predicting genetic parameters and tolerance to pitch canker in slash pine  
– *Andrew Sims*.....pg. 32
- Hyperspectral imaging for the prediction of freeze damage and minimum winter temperature at seed source origin of loblolly pine seedlings  
– *Kitt G. Payn*.....pg. 33

# Rapid-phenotyping of fusiform rust disease resistance in loblolly pine through vibrational spectroscopy

---

**Simone Lim-Hing**<sup>1</sup>, Cristian Montes<sup>2</sup>, Kitt G. Payn<sup>3</sup>, Caterina Villari<sup>4</sup>

<sup>1</sup>Graduate Research Assistant, University of Georgia, Athens, GA, USA, 30606  
(simone.zlim@uga.edu)

<sup>2</sup>Associate Professor, University of Georgia, Athens, GA, USA, 30606

<sup>3</sup>Assistant Professor, North Carolina State University, Raleigh, NC, USA, 27695

<sup>4</sup>Assistant Professor, University of Georgia, Athens, GA, USA, 30606

Fusiform rust is one of the most economically devastating diseases in commercial pine stands in the southeast, causing massive losses in product and revenue. Significant progress has been made in selecting for rust-resistant families; however, the current method of determining fusiform rust disease resistance (i.e. phenotype) at the family level requires lengthy progeny screening trials, which usually need high levels of disease incidence to be effective. Further, this selection process does not directly test the recruitment population, but their progeny only. Thus, a more expedited approach to selection is needed. Here, we propose to use vibrational spectroscopy as an approach to real-time phenotyping for the selection of fusiform rust-resistance in loblolly pine seedlings. Vibrational spectroscopy tools such as Fourier-transform infrared (FT-IR) and near-infrared (NIR) spectroscopy allow users to obtain a comprehensive chemical fingerprint based on the bending and stretching of chemical constituents in a sample. Because pine trees mostly rely on chemical-based defense mechanisms against pathogens, the relationship between chemical makeup and resistance phenotype is promising. In this project, we are going to use a handheld NIR spectrometer to obtain chemical fingerprints from a common set of 40 different families (20 with higher rust incidence and 20 with lower rust incidence) originating from different geographic areas, sampling two non-infected trees per family at five different progeny test sites of the NC State University Cooperative Tree Improvement Program. Collected spectra will be analyzed through machine learning algorithms to build a predictive model that can discriminate the phenotype based on the chemical profile (i.e. chemotype) of a given tree. This project will establish a real-time phenotyping approach that can be used in the field to assess the recruitment population directly, regardless of the disease incidence in the area and without heavily relying on inoculation trials.

# NLR diversity and candidate fusiform rust resistance genes in loblolly pine

---

**Daniel Ence**<sup>1</sup>, Katherine E. Smith<sup>2</sup>, Leandro Gomide Neves<sup>3</sup>, Robin Paul<sup>4</sup>, Jill Wegrzyn<sup>5</sup>, Matias Kirst<sup>6</sup>, Jeremy Brawner<sup>7</sup>, Gary Peter<sup>8</sup>, C. Dana Nelson<sup>9</sup>, and John M. Davis<sup>10</sup>

<sup>1</sup> Postdoctoral Associate, University of Florida, Gainesville, FL, USA 32611 (d.ence@ufl.edu)

<sup>2</sup> Biological Science Technician, Southern Institute of Forest Genetics, Southern Research Station, USDA Forest Service, Saucier, MS, USA, 39574

<sup>3</sup> Chief Executive Officer, Rapid Genomics, Gainesville, FL, USA 32601

<sup>4</sup> Bioinformatics Research Scientist, St. Jude Children's Research Hospital, Memphis, TN, USA 38105

<sup>5</sup> Associate Professor, University of Connecticut, Storrs, CT, USA 06269

<sup>6</sup> Professor, University of Florida, Gainesville, FL, USA 32611

<sup>7</sup> Assistant Professor, University of Florida, Gainesville, FL, USA 32611

<sup>8</sup> Professor, University of Florida, Gainesville, FL, USA 32611

<sup>9</sup> Research Geneticist/Project Leader, USDA Forest Service, Lexington, KY, USA 40546

<sup>10</sup> Senior Associate Dean for Research, IFAS/University of Florida, Gainesville, FL, USA 32611 (jmdavis@ufl.edu)

Fusiform rust causes substantial mortality and productivity losses in pine plantation yields across the southeastern United States. Breeding for genetically resistant families has reduced rust infection. Pine control resistance to the fusiform rust pathogen, *Cronartium quercuum* f.sp. *fusiforme*, in a classic gene-for-gene system. In gene-for-gene interactions, disease symptom expression is conditioned by pathotype-specific genetic interactions between resistance gene alleles and pathogen genotypes harboring specific (a)virulence alleles. Early resistance gene mapping in the loblolly pine (*Pinus taeda*) family 10-5 identified markers for a major fusiform rust resistance gene, *Fr1* (Wilcox et al. 1996; Kuhlman and Powers 1988). More recent work identified markers associated with resistance that mapped to a full-length gene model in version 1.01 of the loblolly pine genome encoding for an NLR protein (Neale et al. 2014). In plants, NLR proteins play key roles in disease resistance to biotrophic pathogens. Given the importance role of NLRs in gene-for-gene disease resistance in model and crop species, their identification and diversity are key targets of research directed at mining in other plant genomes. We hypothesize that elite loblolly pine families selected for disease-resistance and other desirable traits possess novel NLR genes that are not present in the reference genome and that a sequencing strategy targeting novel NLR genes will identify SNP alleles associated with disease resistance. We combined transcriptome mining with sequencing of targeted loci to identify novel NLR genes and map SNPs that reliably segregate with fusiform rust resistance in half-sibling loblolly pine progeny. The novel NLR genes identified here are the first characterization of intraspecies NLRs diversity in a gymnosperm. The SNPs segregating with rust resistance and in novel NLR genes derived from transcriptomes, can be applied to the breeding and deployment of resistant pine seedlings.

# Longitudinal study on in-field trial rust resistance and seedling screening results

---

**Stephanie K. Fong**<sup>1</sup>, W. Patrick Cumbie<sup>1</sup>, Jeremy Brawner<sup>2</sup>, and Michael Cunningham<sup>1</sup>

<sup>1</sup> ArborGen Inc., Ridgeville, SC, USA 29472 (skfong@arborgen.com)

<sup>2</sup> Assistant Professor, University of Florida, Gainesville, FL, USA 32611

The identification and selection of disease-resistant seedlings continues to be an important component of loblolly pine (*Pinus taeda* L.) breeding programs. Fusiform rust (caused by *Cronartium quercuum* (Berk.) Miyabe ex Shirai f.sp. *fusiforme*) is a prevalent disease across much of the southeastern United States and causes significant damage to loblolly pine plantations. Tree breeders can currently evaluate rust resistance through progeny testing in field trials or greenhouse inoculations at the USDA Forest Service Resistance Screening Center (RSC) in Asheville, NC. Both methods to evaluate rust resistance have been useful in breeding programs but the relationship between the two is not always clear. This study explores the relationships between a decade of field trial data and greenhouse inoculation results from the ArborGen population. We will compare how families perform for rust resistance across sites and determine if greenhouse inoculation results correlate to different rust incidence levels in the field. A total of 28 trials were analyzed for rust incidence with individual sites ranging from 4% to 72%. Individual families range from 0% to 80% in the field and 0% to 98% at the RSC.

# Predicting genetic parameters and tolerance to pitch canker in slash pine

---

**Andrew D Sims**<sup>1</sup>, Gary F Peter<sup>2</sup>, and Jeremy T Brawner<sup>3</sup>

<sup>1</sup> Program Manager, Cooperative Forest Genetics Research Program, University of Florida, Gainesville, FL, USA 32611 (adsims@ufl.edu);

<sup>2</sup> Professor, Forest Genetics and Genomics, University of Florida, Gainesville, FL, USA 32611

<sup>3</sup> Assistant Professor, Forest Pathology, University of Florida, Gainesville, FL, USA 32611

Pitch canker is an increasingly important threat to Southeastern forests, especially in slash pine (*Pinus elliottii* var. *elliottii* Engelm.). Its increasing prevalence in the region due to mechanical damage from increased storm activity has led to concern from private and public stakeholders about poor characterization of tolerance from available germplasm. Traditional testing methodologies established to characterize the genetic mechanisms controlling pitch canker tolerance were limited by poor characterization of *Fusarium* isolate virulence and experiment design limitations. Here we describe an alternative experiment design intended to greatly expand the numbers of families that may be screened and increase the power of differentiating among pine families and pathogen isolates evaluated at the USFS Resistance Screening Center. Using this alternative design, genetic parameter estimates for pitch canker tolerance are presented and their implications for the future development of a deployment-oriented slash pine population are discussed.

# Hyperspectral imaging for the prediction of freeze damage and minimum winter temperature at seed source origin of loblolly pine seedlings

---

Yuzhen Lu <sup>1</sup>, Trevor D. Walker <sup>2</sup>, Juan J. Acosta <sup>2</sup>, Sierra Young <sup>3</sup>, Piyush Pandey <sup>2&3</sup>, Austin J. Heine <sup>2</sup>, and **Kitt G. Payn** <sup>2</sup>

<sup>1</sup> Department of Agricultural and Biological Engineering, Mississippi State University, Mississippi State, MS 39762, USA (yzlu@abe.msstate.edu);

<sup>2</sup> Department of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC 27695, USA (kgpayn@ncsu.edu);

<sup>3</sup> Department of Biological and Agricultural Engineering, North Carolina State University, Raleigh, NC 27695, USA

Freeze tolerance is the most important adaptability trait for deployment of loblolly pine in the southeastern US. Minimum winter temperature (MWT) at the seed source origin is the standard indicator of cold adaptability. In this study, we investigated a novel approach for the assessment of freeze-induced damage and prediction of MWT at seed source origin using hyperspectral imaging of seedlings. A population comprising 98 seedlots from Virginia to Florida was subjected to an artificial freeze. A custom-assembled hyperspectral imaging system was used for scanning the seedlings prior to the freeze and on four subsequent occasions after the freeze. On day 44 post the freeze event, seedlings freeze damage was visually scored and the logit link function was used to model the probability of freeze damage. The hyperspectral image data comprised spectra of each seedling that were then averaged for each family, and used to develop family-level predictive models for each scanning date. A significant positive relationship ( $R^2 = 0.28$ ;  $p < .001$ ) between the family MWT and logit scores for freeze damage was observed. Prediction accuracies of freeze damage and MWT based on hyperspectral data varied among scanning dates. The highest prediction accuracy of freeze damage ( $R^2 = 0.79$ ) was achieved using hyperspectral data obtained 41 days after the freeze event. The highest prediction accuracy of MWT ( $R^2 = 0.78$ ) was achieved using hyperspectral data obtained prior to the freeze event. In our study, the best hyperspectral model gave predictions of MWT with a root mean squared error of 2.45 °F, which suggests sufficient precision to rate families for seed source transfer well within a 10 °F plant hardiness zone. Therefore, this research demonstrates that hyperspectral imaging has the potential to serve as a rapid, nondestructive and objective tool for the prediction of MWT of origin of loblolly pine seedlings.



## Session 4: *Lightning Presentations*

---

Genomic assembly comparisons of four Florida *Fusarium circinatum* isolates  
– Katherine Smith.....pg. 36

Genomic characterization of growth and drought tolerance in east Texas loblolly pine  
– Ian MacLachlan.....pg. 37

Can sucrose pathway manipulation affect regrowth of coppiced *Populus*?  
– Trevor Tuma.....pg. 38

A delve into the unknown of sulfate transporters' role in elemental movement and stress response in poplar  
– Samantha Surber.....pg. 39

In search of sassafras with resistance to laurel wilt disease  
– Tyler Dreaden.....pg. 40

Recovering two elite hybrid sweetgum clones for propagation by somatic embryogenesis  
– Nicole A. Locke.....pg. 41

Propagation and conservation of three rare North American ash species using somatic embryogenesis  
– Mason M.W. Richins.....pg. 42

Fusiform rust resistance of loblolly pine inter-provenance hybrids  
– Stephen Goodfellow.....pg. 43

Combined effect of temperature and water stress on germination and plant growth of <i>Magnolia pugana</i> , <i>Populus luziarum</i> and <i>P. primaveralepensis</i> endemic and endangered species from western Mexico – César Jacobo-Pereira.....	pg. 44
Searching orphan genes in hybrid poplar – Chen Hsieh.....	pg. 45
Additive and dominance genetic parameters of mature-aged traits in Arkansas populations of shortleaf pine ( <i>Pinus echinata</i> Mill.) – Chen Ding.....	pg. 46
Conservation genetics and resistance breeding at the Harrison Experimental Forest – Charles Burdine.....	pg. 48
Sustainability genetics for American white oak ( <i>Quercus alba</i> ) – C. Dana Nelson.....	pg. 49
Introduction, evaluation and breeding of novel <i>Populus</i> for subtropical climates – Jason A. Smith.....	pg. 50
Genetic testing based on pedigree reconstruction and spatial analysis in a plantation of <i>Larix kaempferi</i> – Kyungmi Lee.....	pg. 51

# Genomic assembly comparisons of four Florida *Fusarium circinatum* isolates

---

**Katherine E. Smith**<sup>1</sup>, James C. Fulton<sup>2</sup>, Jose C. Huguet-Tapia<sup>3</sup>, Tania Quesada<sup>4</sup>,  
Jeremy Brawner<sup>5</sup>

<sup>1</sup> Biological Science Technician, Southern Institute of Forest Genetics, Southern Research Station, USDA Forest Service, Saucier, MS, USA, 39574 (smithk@ufl.edu);

<sup>2</sup> Postdoctoral Associate, <sup>3</sup> Assistant Scientist, <sup>5</sup> Assistant Professor, Department of Plant Pathology, University of Florida, Gainesville, FL, USA 32611;

<sup>3</sup> Assistant Research Scientist, Forest, Fisheries, and Geomatics Sciences, University of Florida, Gainesville, FL, USA 32611

Pitch canker disease incited by the fungus *Fusarium circinatum*, causes economic losses to the pine timber industry all over the world. Screening seedlings by measuring lesion length following controlled inoculation is used to provide estimates of pine tolerance and isolate virulence. Four isolates that demonstrated differences in virulence when screening slash and loblolly seedlings were selected for further study. To understand the genetic basis for the differences, we used Oxford Nanopore long-read sequencing combined with Illumina short-read sequencing to achieve high quality, nearly chromosome level, genome assemblies. The assemblies were checked for completeness using BUSCO (Benchmarking Universal Single-Copy Orthologs) and aligned to the publically available reference genome (NCBI: GCA\_000497325.3). *Fusarium* species are known to transfer pathogenicity genes and whole “pathogenicity chromosomes” between and within species. Variation in virulence may derive from gene content as well as structural differences in the genome. We located genes in the assembled genomes using annotation software and employed various other techniques to take a closer look at which genes differ between isolates. In addition, we examined the extent of structural differences among the genomes by locating and quantifying the number of sequence inserts, deletions and rearrangements across the genome. This study has identified genomic differences among *Fusarium circinatum* isolates that may be useful for characterizing isolate virulence, monitoring pathogen populations and identifying where different resistance mechanisms should be deployed across the landscape.

# Genomic characterization of growth and drought tolerance in east Texas loblolly pine

---

***Ian R. MacLachlan***<sup>1</sup>, Jason B. West<sup>1</sup>, Casola C<sup>1</sup> and Carol A. Loopstra<sup>1</sup>

<sup>1</sup> Department for Ecology and Conservation Biology, Texas A&M University, 534 John Kimbrough Blvd., College Station, TX 77843 (c-loopstra@tamu.edu)

The loblolly pine populations of east Texas are at the species' warmest range limit. Under future climate warming scenarios the health and productivity of these populations is expected to be negatively impacted. Forestry in east Texas generates billions of dollars in annual revenue and even small negative impacts on annual productivity will have substantial short and long-term economic consequences. Therefore, our research has two primary objectives in context of the east Texas loblolly pine breeding programs. Firstly, to assess the contemporary phenotypic and genomic variation in growth and drought-related traits, and secondly, to identify productive families or trees that are also candidates for increased drought tolerance and future breeding to maintain or increase timber yields. Our approach uses the growth data of 733 trees from 50 maternal families growing in a five-year old progeny trial near Livingston, TX. From these trees we have obtained stable carbon isotope ratios ( $\delta^{13}\text{C}$ ) and nitrogen concentrations, as traits with established relationships to water use efficiency and photosynthetic capacity. In combination with these phenotypic data we are using genotypes from the recently developed Thermo Fisher Axiom™ Pita50K loblolly pine array, and 5000 custom targeted genotype-by-sequencing (GBS) SNPs from drought-related transcripts identified in our previous research. For the GBS we are working with Tecan Genomics to develop a repeatable protocol for using their multiplexed Allegro library preparation platform that has potential to substantially reduce the cost of targeted GBS in conifers. Our preliminary results indicate there is significant variation among families for growth and  $\delta^{13}\text{C}$ , and that array genotyping has been highly successful, yielding ~32,000 high-quality SNPs. More detailed quantitative genetic and association genetic analyses to dissect variation among families and individuals are forthcoming.

# Can sucrose pathway manipulation affect regrowth of coppiced *Populus*?

---

**Trevor T. Tuma**<sup>1</sup>, Batbayar Nyamdari<sup>2</sup>, Scott A. Harding<sup>3</sup>, and Chung-Jui Tsai<sup>4</sup>

<sup>1</sup> PhD Candidate, University of Georgia, Athens, GA, USA 30602 (trevor.tuma@uga.edu);

<sup>2</sup> Research Professional, University of Georgia, Athens, GA, USA 30602

<sup>3</sup> Senior Research Scientist, University of Georgia, Athens, GA, USA 30602

<sup>4</sup> Professor & Georgia Research Alliance Eminent Scholar, University of Georgia, Athens, GA, USA 30602

Perennial woody trees grown for short-rotation bioenergy production are often maintained by coppicing. While it is presumed that non-structural carbohydrate reserves contribute to regrowth, mechanisms for orchestrating remobilization of stored carbohydrates remain poorly understood. Sucrose transport proteins (SUTs) are known for their role in mediating the subcellular mobilization of sucrose from source leaves, yet their involvement in other physiological processes, such as carbohydrate remobilization from sink tissues, remains unclear. Here, we targeted *PtaSUT4* for CRISPR/Cas9-mediated mutagenesis to generate transgenic knock-out (KO) nulls in the *Populus tremula x alba* (INRA 717-1B4) hybrid clone. Amplicon-sequencing confirmed biallelic frameshift mutations in >30 independent transgenic lines; a subset of which was selected for further functional characterization. *SUT4*-KO had no obvious effect on mature plant appearance or growth, but subtle changes to biomass partitioning and metabolism were observed. We then coppiced mature greenhouse-grown poplar to perturb sink-source relations and examine how *SUT4*-KO impacts new vegetative growth. The KO mutants initially exhibited accelerated epicormic bud growth from the stool, but ultimately displayed a decline in new aboveground biomass accumulation after repeated rounds of collection. Metabolic profiling of stool bark and wood during bud flush and regrowth revealed a major shift in primary and secondary metabolism of the mutants. Consistent with vacuolar sequestration in the mutants, sucrose, hexoses, and phenylpropanoid compounds were detected at reduced levels following coppicing and initial bud growth in the mutants. In contrast, the *sut4* mutants accrued elevated levels of amino acids and Krebs cycle intermediates during initial bud burst. This data supports a scenario where carbon was diverted towards primary metabolism to fuel enhanced bud growth in the *sut4* mutants. Collectively, these findings point to a novel role for SUT4 in mobilizing soluble carbohydrates following coppicing and regrowth in *Populus*.

## A delve into the unknown of sulfate transporters' role in elemental movement and stress response in poplar

---

**Samantha Surber**<sup>1</sup>, William P. Bewg<sup>2</sup>, Margot Chen<sup>2</sup>, and Chung-Jui Tsai<sup>1,2,3</sup>

<sup>1</sup> Department of Plant Biology, University of Georgia, Athens, GA, USA 30602  
(sms07287@uga.edu);

<sup>2</sup> Warnell School of Forestry and Natural Resources, University of Georgia, Athens, GA, USA  
30602;

<sup>3</sup> Department of Genetics, University of Georgia, Athens, GA, USA 30602

Sulfur is one of the top six macronutrients essential for plant survival. Lacking this element, plants develop a yellow appearance and declines in biomass. Sulfate is the inorganic form of sulfur that is taken up by the roots and moved throughout the plant to the leaves where it is then assimilated into amino acids, lipids, and other metabolites. Sulfate movement throughout the plant is facilitated by transmembrane proteins known as sulfate transporters (SULTRs). The *SULTR* gene family is split into four groups. Group 1 are high affinity transporters involved in uptake of sulfate from the soil, group 2 are low affinity transporters that interact with group 1 and move the sulfate from the roots to the shoots, group 4 are tonoplast SULTRs for vacuolar efflux, and finally group 3 are the most diverse and lack a cohesive localization and function. Group 3 is made up of five genes, *SULTR3;1*, *SULTR3;2*, *SULTR3;3*, *SULTR3;4*, and *SULTR3;5*. *SULTR3;1* and *SULTR3;5* have been well studied. *SULTR3;1* interacts with the chloroplast and is integral for assimilation of compounds. *SULTR3;5* acts much like groups 1 and 2 in the movement of sulfur from roots to shoots. The other three proteins have conflicting subcellular localizations to the chloroplast or the plasma membrane in the literature. *SULTR3;4* was recently recategorized to be a phosphate transporter in *Arabidopsis* and is phylogenetically related to *SULTR3;3*. This could indicate functional divergence within this subgroup of the SULTR family. In *Populus*, *SULTR3;2* and *SULTR3;4* are highly expressed in the xylem but had opposite sensitivities to development and stressors. CRISPR/Cas9 technology was employed to produce genetic knockouts of both *SULTR3;2* and *SULTR3;4*. These will be subjected to media stress experiments and observed for phenotypic changes such as yellowing. They will also be sent for elemental analysis to see metabolic shifts.

## In search of sassafras with resistance to laurel wilt disease

---

***Tyler J. Dreaden***<sup>1</sup>

<sup>1</sup> Research Plant Pathologist, Southern Research Station, USDA Forest Service, Lexington, KY, USA 40503 (tyler.j.dreaden@usda.gov)

Laurel wilt disease has killed over 300 million redbay in the southeastern USA and is now spreading in sassafras as far north as Kentucky and Tennessee. Currently few if any options are available for managing laurel wilt disease and reducing its effects on forest health. Fungicide injections have been found effective but their costs and need for reapplying make them unsuitable to forests. Cost effective and sustainable options are needed if we are to maintain laurel wilt susceptible hosts as functioning parts of ecosystems. Variation in susceptibility has been identified in redbay and developing host resistance in this species is a promising management option. A collaborative team from the USDA Forest Service, Kentucky Division of Forestry, University of Kentucky, and University of Florida are developing the criteria (e.g., what level of mortality, time since LWD infection, survivorship of hosts in certain diameter classes...) for defining large surviving sassafras from high mortality stands that are likely to have some resistance to laurel wilt disease and not escapes. Using the tool/criteria, we will identify and propagate putatively resistant selections that can be used to develop resistance screening methods and identification of host resistance. Propagated material will be maintained by the University of Kentucky and SRS until disease resistance screening. The tool/criteria will allow others to make their own selections of putatively resistant sassafras from across the region. Ultimately this work will be the start of developing a sustainable and long-term management option using host resistance to maintain sassafras as a functioning member of forested ecosystems.

# Recovering two elite hybrid sweetgum clones for propagation by somatic embryogenesis

---

**Nicole A. Locke**<sup>1</sup>, Paul M. Montello<sup>2</sup>, and Scott A. Merkle<sup>3</sup>

<sup>1</sup> Graduate Research Assistant, University of Georgia, Athens, GA, USA 30602  
(Nicole.Locke@uga.edu);

<sup>2</sup> Research Professional, University of Georgia, Athens, GA, USA 30602;

<sup>3</sup> Professor, University of Georgia, Athens, GA, USA 30602

Somatic embryogenesis-derived hybrid sweetgum (*Liquidambar styraciflua* x *Liquidambar formosana*) clones were developed for pulp and paper and biomass energy applications over 15 years ago. Some of the clones displayed outstanding growth rates and enhanced wood density in field tests. These were propagated as rooted cuttings by ArborGen Inc. and grown by a small number of landowners in the southeastern US, with promising results. However, the clones are currently not being sold by ArborGen. Other clones that were not licensed by ArborGen have been grown by UGA in a small demonstration planting in Athens, GA, for the past 12 years and some of these have demonstrated superior growth rates. With help from consultants, we chose two of the most promising clones to be propagated for further field testing and possible commercialization. Only one of the two clones was still available as an embryogenic culture. Although a copy of this culture that had been maintained continuously by serial transfer no longer was capable of making somatic embryos, two copies of the same culture recovered following 15 years in cryostorage retained this ability. Trees of the other clone in the test planting that was not represented in cryostorage began producing staminate inflorescences in 2018. We collected dormant buds containing inflorescences from this clone in February 2020 and staminate inflorescence tissues were induced to initiate embryogenic cultures at a rate of 13.3%. Of the three plant growth regular treatments tested for induction, only explant material cultured on the basal medium supplemented with NAA produced somatic embryos. Somatic embryos produced from both clones have been germinated to produce somatic seedlings that can be used as the basis for scaled-up propagation of the clones.

# Propagation and conservation of three rare North American ash species using somatic embryogenesis

---

**Mason W.M. Richins**<sup>1</sup>, A. Ryan Tull<sup>2</sup>, and Scott A. Merkle<sup>3</sup>

<sup>1</sup> Graduate Research Assistant, University of Georgia, Athens, GA 30602 USA  
(mwr79514@uga.edu);

<sup>2</sup> Research Professional, University of Georgia, Athens, GA 30602 USA;

<sup>3</sup> Professor, University of Georgia, Athens, GA 30602 USA

In addition to threatening North American ash species of major ecological and economic importance, such as green, white, and black ashes, emerald ash borer (EAB; *Agilus planipennis*) also threatens other native ash trees, of which there are at least 18 North American species. Some of these ash species are comparatively rare. Although not much is known about the susceptibility of most of these species to EAB, some may be at higher risk of extirpation from their native ranges than the more common ash species, due to their smaller ranges and population sizes. Thus, there is a need to extend germplasm conservation efforts to these rare ash species. One approach is to establish new embryogenic cultures of rare ash species for which somatic embryogenesis (SE) has not yet been reported, with the goals using the cultures for propagation and germplasm conservation via cryostorage. We worked with cooperators to locate trees and collect seeds from three rare ash species, Carolina ash (*Fraxinus caroliniana*), Texas ash (*F. albicans*) and Mexican ash (*F. berlandieriana*), for use as explants for SE culture initiation. Immature seeds were harvested from source trees during late summers of 2019 and 2020, and immature zygotic embryos were dissected from them and cultured on media with varying concentrations of 2,4-dichlorophenoxyacetic acid (2,4-D) or picloram. SE was achieved in all three species, with the highest induction rate in Texas ash (12%). Carolina ash somatic embryos from cultures initiated in 2019 were successfully converted to somatic seedlings and hardened off to greenhouse conditions. Somatic embryos from the Texas ash and Mexican ash cultures initiated in 2020 are expected to produce plantlets following pre-germination cold treatment.

## Fusiform rust resistance of loblolly pine inter-provenance hybrids

---

**Stephen Goodfellow**<sup>1</sup>, Adam Polinko<sup>2</sup>, Krishna Poudel<sup>2</sup>, and Phil Dougherty<sup>3</sup>

<sup>1</sup>Graduate Student, Mississippi State University, MS, USA 39762 (sg1942@msstate.edu);

<sup>2</sup>Assistant Professor, College of Forest Resources, Mississippi State University, MS, USA 39762;

<sup>3</sup>Director CFRAM, Dougherty & Dougherty Forestry. Danielsville, GA, USA 30633

Fusiform rust (*Cronartium quercuum* [Berk.] Miyabe ex Shirai f sp. fusiforme) is a noxious pathogen that causes decreased productivity and significant economic loss to forest investments in the southeastern United States. Resistance to fusiform rust is a considerable factor in loblolly pine planting stock selection, particularly in areas with moderate to high rust hazard. Rust resistance has traditionally been evaluated using field trials as well as controlled inoculation trials from the USDA Resistance Screening Center in Asheville, North Carolina. Controlled inoculation screening offers an efficient method of rapidly identifying rust resistant families. Historically, rust resistance research has largely been geographically isolated within provenance. An inter-provenance mating strategy can incorporate rust resistance as well as geographic adaptability of loblolly pine. However, the relative resistance of inter- hybrids across the range remains unclear. This study examines rust resistance of select elite full-sib inter-provenance families, intro-provenance full-sib families and open-pollinated families from parental selections within the Piedmont, Coastal, and Western Gulf geographic origins. We evaluated rust resistance using controlled screening trials and discuss the resistance and deployment of elite inter-province hybrids across the range of loblolly pine.

# Combined effect of temperature and water stress on germination and plant growth of *Magnolia pugana*, *Populus luziarum* and *P. primaveralepensis* endemic and endangered species from western Mexico

---

**César Jacobo-Pereira**<sup>1</sup>, Miguel Ángel Muñiz-Castro<sup>2</sup>, J. Antonio Vázquez-García<sup>2</sup>, Joel Flores<sup>3</sup>, Alejandro Muñoz-Urías<sup>4</sup> and Francisco Martín Huerta-Martínez<sup>4</sup>

<sup>1</sup> Doctorado en Ciencias en Biosistemática, Ecología y Manejo de Recursos Naturales y Agrícolas, Universidad de Guadalajara, Camino Ing. Ramón Padilla Sánchez 2000, 45101 Zapopan, Jalisco, México ( bio.cesarjacobop@gmail.com).

<sup>2</sup> Instituto de Botánica, Departamento de Botánica y Zoología, Centro Universitario de Ciencias Biológicas y Agropecuarias, Universidad de Guadalajara, Camino Ing. Ramón Padilla Sánchez 2000, 45101 Zapopan, Jalisco, México

<sup>3</sup> IPICYT - División de Ciencias Ambientales, Camino a la Presa San José 2055, Col. Lomas 4<sup>a</sup>. Sección, 78216 San Luis Potosí, S.L.P. México

<sup>4</sup> Departamento de Ecología, Centro Universitario de Ciencias Biológicas y Agropecuarias, Universidad de Guadalajara, Camino Ing. Ramón Padilla Sánchez 2000, 45101 Zapopan, Jalisco, México

Since 2016 we have conducted research on the conservation and germination of *Magnolia pugana*, *Populus luziarum* and *P. primaveralepensis* seeds. Our main objective is to know the response in germination and plant growth with the combination of high temperatures and water stress as these conditions are projected for the area where they are distributed. Since 2018 this research is part of the doctoral project of the student César Jacobo Pereira. Germination tests for *M. pugana* have been performed with the combination with three temperatures (24, 28 and 37°C) and five water potentials (0, -0.3, -0.6, -0.9 and -1.2 MPa). The interaction of temperature and water potential on germination percentages was significant (ANOVA:  $F = 3.86$ ,  $P < 0.001$ ). The highest number of germinated seeds (78%) was obtained at a temperature of 24°C and 0 MPa. These results indicate that germination of *M. pugana* is severely affected by the interaction between increasing temperatures and decreasing precipitation. In *P. luziarum* and *P. primaveralepensis* we tested germination and storage of their seeds at two temperatures. In the first 24 hrs after seed collection, germination was high (91 and 95%, respectively). Germination percentages decreased when stored at 21°C and for more than four weeks. The germination percentage of *P. primaveralepensis* decreased more slowly than that of *P. luziarum* at 4°C. This indicates that subtropical *Populus* subtropical seed storage conditions respond similarly to those in temperate climates. If approved, the grant for which I want to participate will serve to continue my studies and experiments to generate crucial information to establish in situ and ex situ conservation and repopulation programs.

## Searching orphan genes in hybrid poplar

---

**Chen Hsieh**<sup>1</sup>, Ran Zhou<sup>2</sup>, and Chung-Jui Tsai<sup>3</sup>

<sup>1</sup> PhD student, Institute of Bioinformatics, University of Georgia, Athens, GA 30602, USA  
(Chen.Hsieh@uga.edu);

<sup>2</sup> Post-Doctoral Associate, Warnell School of Forestry and Natural Resources, University of Georgia, Athens, GA 30602, USA;

<sup>3</sup> Professor, School of Forestry and Natural Resources, Department of Plant Biology, Department of Genetics and Institute of Bioinformatics, University of Georgia, Athens, GA 30602, USA

Orphan genes by definition have restricted taxonomic occurrence. They don't share sequence similarity to other known genes in the bioinformatic databases, thus making it difficult to predict their biological function using traditional approaches. Since orphan genes are unique to specific taxa, they may confer adaptive significance during evolution. We focus on the *Populus tremula* × *P. alba* INRA 717-1B4 hybrid (hereafter '717'), an important study system in poplar functional genomics because of its high transformation efficiency. To identify protein-coding orphan genes, a total of 15 genomes were selected, including well annotated model species and those closely related to our focal species for analysis. Orthogroups were constructed according to the amino acid sequence similarity among the selected proteomes. Genes belonging to orthogroups that only contain *Populus* sequences as well as those which cannot be assigned to any orthogroups were categorized as putative orphan candidates. Collinearity of certain 717-specific orphan genes can be observed between the two subgenomes in the current 717 draft genome assembly. The candidate orphans showed several properties, including shorter coding sequences and higher isoelectric points. Among these orphan genes, seven showed interesting expression patterns in different tissues and under drought stress treatment. Three genes showed higher expression in xylem. Two genes were only expressed in callus and the predicted polypeptides are shorter than 100 amino acids. The other two genes were induced under drought stress and were only expressed in roots. In addition, they were all predicted to contain intrinsically disordered regions. We hypothesize that these candidate orphans may act as chaperone proteins and contribute to the fitness of '717'.

## Additive and dominance genetic parameters of mature-aged traits in Arkansas populations of shortleaf pine (*Pinus echinata* Mill.)

---

**Chen Ding**<sup>1</sup>, Shaik Hossain<sup>2</sup>, Virginia McDaniel<sup>3</sup>, Benjamin Rowland<sup>4</sup>, Benjamin Bartlett<sup>5</sup>, Fred Raley<sup>6</sup>, Carolyn Pike<sup>7</sup>, Barbara S. Crane<sup>8</sup>, Don C. Bragg<sup>9</sup>, C. Dana Nelson<sup>10,11</sup>

<sup>1</sup>Geneticist, Western Gulf Forest Tree Improvement program, Texas AM Forest Service, TAMU, College Station, TX 77843, USA, chen.ding@email.tamu.edu

<sup>2</sup> Assistant Professor, Department of Biological & Environmental Sciences, Alabama A&M University, Normal, AL 35762

<sup>3</sup> Ecologist, USDA Forest Service, Southern Research Station, Hot Springs, AR 71901

<sup>4</sup> Seed Orchard Manager, USDA Forest Service, Southern Region, Mt. Ida, AR 71957

<sup>5</sup> Tree Improvement Coordinator, Western Gulf Forest Tree Improvement program, Texas AM Forest Service, TAMU, College Station, TX 77843, USA

<sup>6</sup> Director, Western Gulf Forest Tree Improvement program, Texas AM Forest Service, TAMU, College Station, TX 77843, USA

<sup>7</sup> Forest Regeneration Specialist, USDA Forest Service, Eastern Region, State and Private Forestry, West Lafayette, IN

<sup>8</sup> Regional Geneticist (retired), USDA Forest Service, Southern Region, Atlanta, GA

<sup>9</sup> Research Forester, USDA Forest Service, Southern Research Station, Monticello, AR

<sup>10</sup> Research Geneticist, USDA Forest Service, Southern Research Station, Forest Health Research and Education Center, Lexington, KY

<sup>11</sup> Project Leader, SDA Forest Service, Southern Research Station, Southern Institute of Forest Genetics Saucier, MS 39574, USA, charles.d.nelson@usda.gov

Range-wide ecosystem restoration efforts are in progress for shortleaf pine (*Pinus echinata*). Genetic considerations must include provenance or seed source and genetic diversity as well as performance of the planted seedlings. Genetic information on seed source performance has been documented through provenance tests conducted in the 1950s-1980s. Genetic diversity studies have shown an increase in hybridization with loblolly pine from the 1950s to the 2000s. Only limited information is available on within source variation attributed to individual parents. We remeasured 15 R8 progeny tests representing the Ouachita and Ozark National Forests seed sources. Disconnected half-diallel crosses were made among the first-generation parents in the Mt Ida, AR seed orchard and established at tests sites in both national forests. Earlier measurements (ages 5 and 10 years), available on some of these tests, were merged with the current measurements that ranged from 31 to 40 years from planting. In total, 21,260 planted trees were evaluated for survival through the current assessment age, with all surviving trees measured for DBH and scored for damage and straightness, and a sample of these trees were measured for height. This data set provided progeny test performance information on 126 parents and 330 full-sib families. We found substantial additive and dominance genetic variation in growth and straightness, while the proportions of these variances differed by traits and ages. Additive and dominance

genetic variances increased with age resulting in larger narrow- and broad-sense heritabilities for height at the latest measurement age. Predicted genetic gain is promising to advance tree growth and straightness for reforestation using both open- and control-pollinated families. Moderate age-age correlations suggest selection of high-performing parents in older trials can be done at younger ages.

## Conservation genetics and resistance breeding at the Harrison Experimental Forest

---

**Charles Burdine**<sup>1</sup>, Tyler J. Dreaden<sup>2</sup>, Katherine Smith<sup>3</sup>, Warren Nance<sup>4</sup>, Austin Himes<sup>5</sup>,  
Ellen Crocker<sup>6</sup>, Jeremy Brawner<sup>7</sup>, Jason A. Smith<sup>8</sup>, and C. Dana Nelson<sup>9</sup>

<sup>1</sup> Biologist, USDA Forest Service, Saucier, MS, USA 39574;

<sup>2</sup> Research Plant Pathologist, USDA Forest Service, Lexington, KY, USA 40546;

<sup>3</sup> Biologist, USDA Forest Service, Gainesville, FL, USA 32611;

<sup>4</sup> Research Geneticist (retired), USDA Forest Service, Saucier, MS, USA 39574;

<sup>5</sup> Assistant Professor, Mississippi State University, Starkville, MS, USA 39762;

<sup>6</sup> Assistant Professor, University of Kentucky, Lexington, KY, USA 40546;

<sup>7</sup> Assistant Professor, University of Florida, USA, 32611;

<sup>8</sup> Associate Professor, University of Florida, USA, 32611;

<sup>9</sup> Research Geneticist, USDA Forest Service, Lexington, KY, USA 40546

(charles.d.nelson@usda.gov)

Genetic resistance offers a promising, sustainable option for managing introduced pests and pathogens in forestry. To counter two such introduced pathosystems, chestnut blight and laurel wilt, we are initiating conservation genetics and resistance breeding projects in the host species. Chestnut blight (caused by *Cryphonectria parasitica*) has virtually eliminated American chestnut (*Castanea dentata*) from the landscape since being introduced more than a century ago. We are searching throughout Mississippi for American chestnuts to use in cooperative resistance breeding. Mississippi is a desirable search area, since it represents the most southern portion of the species native range. American chestnuts will be vegetatively propagated, planted in two breeding orchards in Mississippi and genotyped to help validate species identity. To date, 20 putative American chestnuts have been discovered and propagated for orchard planting and genotyping. Laurel wilt (caused by *Raffaelea lauricola*) has killed over 300 million host trees since being discovered in Georgia in 2002. The main host tree has been redbay (*Persea borbonia*) in the southeast, but the disease is now impacting sassafras (*Sassafras albidum*) in areas as far north as Tennessee and Kentucky. Tolerance to the pathogen has been identified in redbay and we will take a similar approach to identify tolerance in sassafras. Specifically, we are propagating putatively tolerant selections of redbay and sassafras and will begin screening them for resistance to identify individuals for our breeding orchard. Currently, eight LW-tolerant redbay selections from the University of Florida are being established in the breeding orchard in Mississippi. A multiagency team is establishing a study to select individual sassafras trees with potential tolerance to laurel wilt. Disease resistance screening will be conducted once selections are propagated. Trees showing levels of resistance will be mated to evaluate the trait's inheritance and to identify candidate trees for breeding and seed orchard development.

## Sustainability genetics for American white oak (*Quercus alba*)

---

**C. Dana Nelson**<sup>1</sup>, Albert G. Abbott<sup>2</sup>, Laura DeWald<sup>3</sup>, Mark V. Coggeshall<sup>4</sup>, Margaret E. Staton<sup>5</sup>,  
Beant Kapoor<sup>5</sup>, Jozsef Stork<sup>6</sup>, Shenghua Fan<sup>6</sup>, Tatyana Zhebentyayeva<sup>7</sup>, Rachel Thunder<sup>3</sup>,  
Nurul Faridi<sup>8</sup>, John E. Carlson<sup>7</sup>, Ellen V. Crocker<sup>3</sup>, John Lhotka<sup>3</sup>, Jian Yang<sup>3</sup>,  
Jeffrey W. Stringer<sup>3</sup>, and Seth DeBolt<sup>9</sup>

<sup>1</sup>USDA Forest Service, Southern Research Station, Lexington, KY (charles.D.Nelson@usda.gov);

<sup>2</sup>Forest Health Research and Education Center, University of Kentucky, Lexington, KY;

<sup>3</sup>Department of Forestry & Natural Resources, University of Kentucky, Lexington, KY;

<sup>4</sup>School of Natural Resources, University of Missouri, Columbia, MO;

<sup>5</sup>Department of Entomology & Plant Pathology, University of Tennessee, Knoxville, TN;

<sup>6</sup>Department of Horticulture, University of Kentucky, Lexington, KY

<sup>7</sup>Department of Ecosystems Sciences & Management, Penn State University, State College, PA;

<sup>8</sup>USDA Forest Service, Southern Research Station, College Station, TX;

<sup>9</sup>Beam Institute for Kentucky Spirits, University of Kentucky, Lexington, KY

White oak (*Q. alba*) is a keystone forest tree species distributed over much of the eastern US. The species is important ecologically for watershed and wildlife health, and economically for lumber, veneer, and specialty products such as barrel staves. Because of the species importance, there are sustainability concerns primarily driven by forest mesophication, limiting regenerated seedlings' recruitment into the canopy, and climate change. The recently formed White Oak Initiative speaks to these concerns and looks to partner on key research themes encompassing white oak's biological potential in forest management. An important determinant in biological potential is a species' genetic diversity and how this diversity might be characterized and utilized in forest management. Our stakeholder-supported research program is addressing white oak genetics in three ways— (1) germplasm collection, conservation, and improvement; (2) genomic, transcriptomic and metabolomic assessment of genetic diversity; and (3) characterization of adaptive genes through analysis of genotype-phenotype associations. Initial work has resulted in two years of range-wide acorn collections, including a first-year nursery crop with subsequent field test plantings, and completion of a diversity study of white oak stands in the Daniel Boone National Forest. We are currently developing a high-quality, white oak genome sequence and comprehensive transcriptome and metabolome resources. Subsequent work will continue range-wide acorn collections and field test plantings, as well as finalizing and annotating the genome and initiating trait phenotyping and genotyping.

# Introduction, evaluation, and breeding of novel *Populus* for subtropical climates

---

**Jason A. Smith**<sup>1</sup>, Nicolas Anger<sup>2</sup>, Jonathan Lubar<sup>3</sup>, Patrick McGovern<sup>4</sup>, Victor Hugo Garza Guajardo<sup>5</sup>, Miguel Ángel Muñiz-Castro<sup>6</sup> and J. Antonio Vázquez-García<sup>7</sup>

<sup>1</sup> Associate Professor, University of Florida, Gainesville, FL, USA 32611 (jasons@ufl.edu)

<sup>2</sup> PhD Candidate, University of Florida, Gainesville, FL, USA 32611

<sup>3</sup> Horticulturist, University of Florida, Gainesville, FL, USA 32611

<sup>4</sup> Private Nurseryman and Middleware Administrator, Grand Rapids, MI 49525

<sup>5</sup> San Nicolás de los Garza, Código Postal 66422, Nuevo León, Mexico

<sup>6</sup> Professor, Herbario IBUG (“Luz María Villarreal de Puga”), Laboratorio de Ecosistemática, Inst. de Botánica, Dpto. de Botánica y Zoología, Univ. de Guadalajara-CUCBA, C. P. 45200, Zapopan, Jalisco, Mexico.

<sup>7</sup> Professor, Herbario IBUG (“Luz María Villarreal de Puga”), Laboratorio de Ecosistemática, Inst. de Botánica, Dpto. de Botánica y Zoología, Univ. de Guadalajara-CUCBA, C. P. 45200, Zapopan, Jalisco, Mexico.

Since 2008 the Forest Pathology Lab at the University of Florida has established and evaluated *Populus* germplasm with potential adaptation to subtropical climates such as Florida. We aimed to examine disease resistance, photoperiod response, growth rate, ease of propagation, adaptability to heat and lack of chilling. To this end, the focus has been to grow species from Mexico, naturally and allopatrically occurring within seven different biogeographic provinces in subtropical low-latitude locations (18°48’-30°34’ N), including *P. aff. fremontii* var. *mesetae* (Jalisco origin), *P. luziarum*, *P. primaveralepis*, *P. mexicana*, *P. guzmanantlensis*, *P. monticola*, *P. simaroa*, *P. tremuloides* (Jalisco origin) and evergreen clones originating from the mutant form of Lombardy poplar, *P. nigra* ‘Chile’, first discovered over 100 years ago near Santiago, Chile as a sport mutation on *P. nigra* ‘Italica’. We carried out progeny trials (100 seedlings each) with *P. mexicana* ssp. *mexicana* using three open-pollinated families from near Monterrey, Mexico. Results from year one (from seed) suggested significant potential with this species, with mean height growth of > 4 m and very high disease and insect pest resistance. Seed propagation for *P. luziarum* and *P. primaveralepis* was highly successful and although height growth for both species after six months was not significantly different, the means were 22.91 cm (± 2.19 cm) and 25.78 cm (± 3.45 cm), respectively. Early flowering was observed for the white poplar *P. luziarum*, with both male and female clones flowering within the first year from cuttings. Pollen was collected from *P. luziarum* and *P. monticola*, and crosses with *P. alba* were attempted, but so far no viable seed has been produced. This is the first report on our observations and early results suggest there is significant potential within these species for trait discovery, adaptation and future development within the genus for warmer climates.

# Genetic testing based on pedigree reconstruction and spatial analysis in a plantation of *Larix kaempferi*

---

**Kyungmi Lee**<sup>1</sup>, Jeong Ho Song<sup>2</sup>, In-Sik Kim<sup>3</sup>, and Kyu-Suk Kang<sup>4</sup>

<sup>1</sup> Researcher, Forest Tree Improvement and Biotechnology Division, Forest Bioresources Department, National Institute of Forest Science, Suwon 16631, Republic of Korea (kmile@korea.kr);

<sup>2</sup> Senior Researcher, Forest Tree Improvement and Biotechnology Division, Forest Bioresources Department, National Institute of Forest Science, Suwon 16631, Republic of Korea;

<sup>3</sup> Director, Forest Tree Improvement and Biotechnology Division, Forest Bioresources Department, National Institute of Forest Science, Suwon 16631, Republic of Korea;

<sup>4</sup> Professor, Department of Agriculture, Forestry and Bioresources, College of Agriculture and Life Sciences, Seoul National University, Seoul 08826, Republic of Korea

*Larix kaempferi* is one of the major timber species in Northeast Asia because of their rapid growth and straightness. The demand for the reforestation of the species has been increased in Korea due to the promoted timber utilization by recent advances in wood processing technology. The genetic testing is the prerequisite process for the evaluation of genetic value and gain, and in turn for the supply of genetically improved seed. However, the genetic testing of *L. kaempferi* has not been performed yet because it is an introduced species in South Korea. The establishment of progeny trial of the species is even difficult because of the irregularity and the large variation in seed production. In this study, the genetic testing was simulated using the pedigree reconstruction and the spatial distribution analysis in a plantation originated from seed orchard crops in order to replace progeny trials. The adequacy of utilizing the plantation as a testing population was confirmed based on the comparable level of the genetic variation in the offspring to that in the group of mother trees. The pedigree reconstruction was conducted by maternity analysis using microsatellite markers. The genetic testing of diameter growth was performed subsequently using both the animal model and the model accounting for spatial autoregression. The improved fitness of the latter presented the usefulness of the spatial analysis in genetic testing with plantation lacking the prior experimental design. To understand the environmental characteristics of the heterogeneous site, the environmental effects on the growth were investigated by geographically weighted regression analysis. The changes in the main effects allowed it possible to propose a zoning scheme for the management of the plantation as breeding material. In addition, it could be expanded to be utilized in the genetic testing of the other tree species without progeny trials.



## Session 5: *Schatz Symposium*

---

- Epigenetic influences on plant growth response, and potential for enhancing stress resilience in tree species  
– *Sally Mackenzie*.....pg. 53
- A quantitative Reduced Representation Sequencing (qRRS) of genomes; a paradigm shift in NGS-based genotyping  
– *Bode Olukolu*.....pg. 54
- Genomics in rare species conservation: Teasing apart evolutionary history and adaptation  
– *Jill Hamilton*.....pg. 55

# Epigenetic influences on plant growth response, and potential for enhancing stress resilience in tree species

---

**Sally A. Mackenzie**<sup>1</sup>, Hardik Kundariya<sup>1</sup>, Roberly Sanchez<sup>1</sup>, Xiaodong Yang<sup>1</sup>, Ha Eun Jeh<sup>2</sup>, Isaac Dopp<sup>2</sup>

<sup>1</sup> Departments of Biology and Plant Science, Penn State University (sam795@psu.edu);

<sup>2</sup> Plant Biology Graduate Program, Penn State University, University Park, PA 16802

Understanding plant environmental response, and the heritability of that response, may prove vital to dissecting modes of plant adaptation. Our group has developed a system for investigating plant epigenetic behaviors that centers on plastid triggers of epigenetic reprogramming. These plastid-based epigenetic effects appear to be well conserved in plants and rely on histone modifying, DNA methylation and small RNA components. Implementation of epigenetic reprogramming in grafting experiments permits the identification of RNA-directed DNA methylation targets within the genome that involve graft transmissible and transgenerational signaling and lead to enhanced fitness phenotype changes. Two of the genetic components of this system, MSH1 and PPD3, were first identified and characterized in Arabidopsis, but are conserved in plants and offer important potential strategies for enhancement of growth potential and resilience in tree species. We have developed a model for plant environmental response that may apply to both seasonal and perennial growth habits, with a plan for implementation in poplar for identifying environmentally altered gene networks and alignment with longer-term genome evolutionary paradigms.

# A quantitative Reduced Representation Sequencing (qRRS) of genomes; a paradigm shift in NGS-based genotyping

---

**Bode A. Olukolu**<sup>1</sup> and G. Craig Yencho<sup>2</sup>

<sup>1</sup> Assistant Professor, Department of Entomology and Plant Pathology, University of Tennessee, Knoxville, TN USA 37996 (bolukolu@utk.edu); <sup>2</sup> Professor, Department of Horticultural Science, NC State University, Raleigh, NC USA 27695.

Next Generation Sequencing (NGS) is an extensively used tool for massive parallel sequencing of genomes. However, several applications such as NGS-based genotyping still suffer from pitfalls that limit their accuracy and utility. To mitigate these pitfalls, we have developed a significant advancement in short-read NGS library preparation. Presented here is an inexpensive quantitative reduced representation sequencing (qRRS) approach for dosage-sensitive genotyping and quantitative strain-level metagenome/microbiome profiling. The scalable, ligation-free and double-stranded DNA-protection assay eliminates off-target annealing temperature-dependent hybridization. This is achieved by using single-stranded barcoded adapters for isothermal strand displacement of double-stranded DNA templates with restriction site overhangs as the only priming site. As much as 9,216 samples can be multiplexed. Novel features in this protocol include a paradigm shift in adapter design that prevents chimeric reads and barcode swapping, a flow cell cluster enhancer that generates about 50% more yields, and consistent high-quality base calling scores. The library preparation workflow is optimized for ease-of-use and can be completed in one to two days. To accommodate these novel features during data pre-processing and downstream analytics, we have developed bioinformatic and analytical pipelines for empirical-based NGS data quality filtering (ngsComposer), haplotype-based variant calling and filtering (GBSapp), and quantitative metagenomic alignment and taxonomic exact matching (Qmatey). The qRRS approach establishes new standards in high-fidelity quantitative genotyping, minimizes missing data and allelic dropout, and makes functional microbiome studies more accessible. Compared to 16S amplicon sequencing, which uses a single gene for microbiome profiling, qRRS provides multiple genome-wide sequences for strain-level taxonomic delineation and quantification. We are now exploring its utility as a diagnostic tool for scoring multiple diseases (and disease complexes) based on titer levels of pathogens in a single assay. We envision that the enhanced quality and quantity of qRRS-derived markers will improve genomic-assisted breeding efforts.

## Genomics in rare species conservation: teasing apart evolutionary history and adaptation

---

**Jill Hamilton**<sup>1</sup>, F. Thomas Ledig<sup>2</sup>, Lionel Di Santo<sup>1</sup>, and Jessica Wright<sup>3</sup>

<sup>1</sup> North Dakota State University, Fargo, ND, USA (jill.hamilton@ndsu.edu);

<sup>2</sup> Department of Plant Science, University of California, Davis, CA, USA;

<sup>3</sup> Pacific Southwest Research Station, USDA-Forest Service, Davis, CA, USA

Species evolutionary potential is tightly linked to both the amount and distribution of genetic variation available through which natural selection may act. Indeed, the genetic consequences of isolation and population size may be exacerbated in rare species, limiting species' ability to adapt to ongoing change. Thus, in a rapidly changing environment, maintenance of genetic variation within and across populations becomes an increasingly important target for species conservation. Here, we discuss the importance of understanding species' evolutionary history, and the role different evolutionary processes may play influencing neutral and adaptive processes both across space and time. We discuss the importance of these data to establishing conservation collections and designing species management strategies that preserve species' evolutionary potential. Providing a case study, we focus on Torrey pine (*Pinus torreyana* Parry) a critically endangered pine endemic to California. The combination of small population size, extremely low genetic variation, and abiotic and biotic challenges associated with climate change indicate Torrey pine may have reduced evolutionary potential to adapt to change. Thus, Torrey pine may be a potential candidate for inter-population genetic rescue. Pairing genomic data with phenotypic data from natural populations and common garden experiments, Torrey pine provides an ideal system to evaluate the contribution of demographic history, gene flow, and natural selection to population differences. These data become essential to considering potential risks associated with management decisions and identifying short and longer-term conservation strategies necessary to preserve species' evolutionary potential.