

# Biological Nutrient Management: Best Organic Practices for Soil Fertility and Resource Stewardship

*A Webinar for NRCS  
February 6th 2023*

Presented by:

Mark Schonbeck, Research Associate, Organic Farming Research Foundation (OFRF)

## Presentation Notes

### Introduction (Slides 1-4)

#### Slide 2 – *Goals of nutrient management*

Production goals include *optimum* yield, taking direct and environmental production costs into consideration; therefore, not necessarily maximizing yield. Conservation concerns focus mainly on the adverse impacts of soluble nitrogen (N) and phosphorus (P) on water quality, and more recently on greenhouse gas (GHG) emissions from fertilized soils.

Organic farmers take a resource-conserving approach to nutrient management to build the soil's nutrient retention, cycling, and delivery capacity, thereby reducing the amount of nutrient input required for satisfactory yield.

#### Slide 3 – *Nutrient management criteria* (NRCS CPS 590 and NOP Standards)

Certified organic producers may also use the following non-biological nutrient sources and pH-correcting amendments (USDA, NOP, undated):

- Natural (mined) mineral substances of low solubility, e.g. calcitic or dolomitic limestone.
- Natural sodium nitrate (“Chilean nitrate”) to provide no more than 20% of the crop’s total N requirement.
- Other natural (mined) mineral substances of high solubility (e.g., potassium sulfate, langbeinite or sulpomag), *except* those on the National List of prohibited non-synthetic substances (potassium and calcium chlorides, lead, arsenic, and fluo-aluminate minerals).
- Elemental sulfur
- Magnesium sulfate for documented Mg deficiency
- Soluble boron compounds for documented B deficiency.
- Sulfates, carbonates, oxides, or silicates of zinc, copper, iron, manganese, molybdenum, selenium, and cobalt when micronutrient deficiency is identified by soil or tissue test.

#### Slide 4 – *Soil fertility: organic farmers’ perspectives*

In 2020, the Organic Farming Research Foundation (OFRF) conducted a survey of over 1,000 certified organic and 71 transitioning-organic producers to document their use of resource stewardship and production practices, their leading production and enterprise management challenges, and technical assistance needs (Snyder et al., 2022).

Our findings illustrate the extent to which organic farmers implement the fertility management practices codified in the USDA National Organic Program (NOP) Standards. For example, 68% of respondents reported using cover crops regularly (“often” or “very often”), a figure which increased to 76% for organic field crop farmers, compared with perhaps 10% of conventional field crop farmers. Nearly three out of four survey respondents have established one or more perennial conservation plantings, which protect water quality by intercepting nutrient runoff or leaching.

Nutrient management and soil health were the second and third most-cited technical assistance needs after organic weed, pest, and disease management.

### Organic Nutrient Management 101 (Slides 5-17)

Evolving perspectives on soil nutrient dynamics and the central role of soil organisms.

#### *Slide 6 – 20<sup>th</sup> Century nutrient management*

With the development of the Haber-Bosch process for converting inert atmospheric nitrogen (N<sub>2</sub>) into ammonia (NH<sub>3</sub>) and other industrial processes for converting insoluble phosphorus (P) and potassium (K) minerals into soluble forms, 20<sup>th</sup> Century farmers found a “quick fix” for declining soil fertility and crop yield, which were attributed to shortages of nitrogen N, P, and K.

During the 1950s-1970s, standard soil test recommendations for NPK applications were based on estimates of crop nutrient uptake, yield goals, and yield response trials conducted on research station soils that had become depleted from years or decades of conventional management with excessive tillage and inadequate return of organic residues to sustain soil life. The community of soil life was mostly overlooked or seen as competing with the crop for nitrogen and other nutrients, tying-up soluble soil N as it decomposes organic residues. Recommended NPK rates were further increased to account for anticipated losses through runoff, leaching, denitrification, microbial immobilization, and mineral-fixation of phosphorus.

The reason to dwell briefly on the past is that this paradigm still informs 21<sup>st</sup> century standard soil test recommendations to a significant degree. On biologically active soils such as those that develop over time under organic management, the assumptions underlying this approach to identifying the right rates and forms of nutrients are out of date.

#### *Slide 7 – 20<sup>th</sup> Century organic farming: organic matter for fertility*

The organic farming movement of the early-mid 20<sup>th</sup> Century raised concerns that the new fertilizer technology could not sustain the health of the land, crops, livestock, or people because it bypasses the natural soil-based processes of recycling organic matter and plant nutrients from plant and animal residues to support new plant life and agricultural production.

Early practitioners of the organic method implemented various “feed the soil” strategies that mimic natural processes and parallel the soil health practices advocated by NRCS over the past 20 years. They also sought to integrate crop and livestock production to improve nutrient and resource cycling, create a more “whole” farm ecosystem, and reduce reliance on off-farm inputs. Synthetic fertilizers and pesticides were excluded to protect soil micro-organisms and earthworms, whose importance to soil fertility they understood.

Slides 8-9 – 20<sup>th</sup> Century organic nutrient management: thinking beyond soluble NPK; micronutrients for crop health and nutritional value

From the beginning of the organic movement, practitioners recognized that soluble forms of NPK are not the only relevant forms of nutrients, that N, P, and some other nutrients exist in organic forms that plants can access provided that the soil is healthy, with an active and well-fed community of life that converts organic and even mineral-bound nutrients into plant-available forms. In addition, organic farmers and agricultural advisors paid attention to the whole spectrum of essential elements including the trace minerals or micronutrients essential to plants, animals, humans, and the nutritional quality of farm products (see Table 1 below). Note that several micronutrients play essential roles in N fixation and uptake of N and P.

Organic practices – crop rotation, cover crops, and using complex organic and natural-mineral fertility sources – promote balanced crop nutrition and help to provide trace elements.

Plant-available anions (negative charge) like nitrate, phosphate, sulfate, and borate exist mainly in the soil solution, and can easily leach out of the root zone during heavy rainfall or snowmelt. Plant available cations (positive charge) like K, calcium (Ca), magnesium (Mg), ammonium nitrogen ( $\text{NH}_4^+$ ), and some micronutrients can also occur in the soil solution but are mostly adsorbed (held) on the soil's cation exchange capacity consisting of negatively charged clays and stable soil organic matter. Cations are less prone to leaching except in sandy soils low in organic matter and clay.

Nutrient reserves (not immediately available to plants, but potentially released through microbial and plant root biological processes) are held in soil organic matter (N, P, S, some micronutrients) and in soil minerals (P, K, Mg, Ca, micronutrients).

Soil organic matter and soil life play important roles in providing essential micronutrients for crops and livestock. Low-solubility cations like copper (Cu), zinc (Zn), iron (Fe), and manganese (Mn) become more plant available when they combine with soluble organic substances in the soil to form chelates, when soil microbes convert them to more soluble forms, and when mycorrhizal fungi facilitate plant uptake. Favorable soil pH (6.0 – 7.0 for most crops, and ~5.0 for blueberry and cranberry) plays an important role in crop access to micronutrients.

Boron is present as a soluble anion, and is often deficient in soils under warm, rainy climates. Sodium (Na) and chlorine (Cl) are rarely deficient and can be present in excess in saline-alkaline soils of semiarid regions, or in overfertilized soils, especially in high tunnels.

Soil tests for micronutrients often do not accurately reflect actual crop nutritional status with regards to these essential trace elements. Depending on the crop and variety, soil-root microbiome, overall soil health, soil mineralogy, and pH, crops may obtain sufficient amounts of micronutrients that shows “low” on the soil test, or conversely suffer deficiency despite a “sufficient” soil test value. Therefore, organic farmers (and many non-organic specialty crop farmers as well) often use a foliar nutrient analysis to assess crop nutritional status and identify micronutrient supplement needs more accurately.

Moderate deficiencies can often be corrected with the use of seaweed or fish-based foliar fertilizers or soil amendments, or compost made from a diversity of materials. More severe or persistent deficiencies can be addressed with NOP-approved micronutrient minerals. Good organic management, maintaining desirable soil pH, compost made from a diverse mix of materials, and additional supplementation when indicated by foliar and soil tests can ensure optimal levels of micronutrients.

Table 1. Essential macro and micronutrients, their forms in soil, and their roles in plant nutrition. Based on information in Weil and Brady (2017).

Element	Forms in the soil <sup>1</sup>	Essential nutritional roles
Nitrogen (N)	Anion, cation, organic	Structural component of proteins, enzymes, amino acids, chlorophyll, nucleic acids, vitamins, secondary compounds
Phosphorus (P)	Anion, organic, mineral	Structural component of nucleic acids, cell membranes (phospholipids), and ATP (“energy currency” of all organisms)
Potassium (K)	Cation, mineral	K <sup>+</sup> in solution vital for photosynthesis, energy production, enzyme activation, stress resilience
Sulfur (S)	Anion, organic, mineral	Structural component of proteins, enzymes, vitamins, secondary compounds
Calcium (Ca)	Cation, mineral	Cell wall component, membrane permeability, enzyme activation
Magnesium (Mg)	Cation, mineral	Central ion in chlorophyll molecule, activation of enzymes; energy, oil, and protein synthesis.
Sodium (Na)	Cation	Essential for animals and humans, can take some but not all of the roles of K <sup>+</sup> in plants
Silicon (Si)	Anion, mineral	Strengthens plant structure, stress resilience. <i>More research is needed and is ongoing.</i>
Iron (Fe)	Cation, chelate, mineral	Structural part of nitrogenase (N fixation), N and P nutrition, chlorophyll synthesis
Manganese (Mn)	Cation, chelate, mineral	Enzymes for photosynthesis and N metabolism
Zinc (Zn)	Cation, chelate, mineral	Multiple enzymes, seed production and maturation
Boron (B)	Soluble boric acid (H <sub>3</sub> BO <sub>3</sub> ), anion in alkaline soils	Enzymes, cell division and development, sugar translocation, nucleic acid and hormone synthesis
Copper (Cu)	Cation, chelate, mineral	Enzymes, photosynthesis, protein and carbohydrate metabolism, N fixation
Nickel (Ni)	Cation, chelate, mineral	Enzymes, seed filling and viability, N fixation
Molybdenum (Mo)	Anion, chelate, mineral	Structural part of nitrogenase (N fixation) and nitrate reductase enzyme
Chlorine (Cl)	Anion	Enzyme activation, photosynthesis
Cobalt (Co)	Cation, chelate, mineral	Essential for N-fixing bacteria, animals, people
Selenium (Se)	Anion, mineral	Essential for animals and people
Chromium (Cr)	Anion, cation, mineral	Possibly essential for animals and people

<sup>1</sup> Anion = negatively charged ion (e.g. NO<sub>3</sub><sup>-</sup>) in soil solution or held on anion exchange sites in soil. Cation = positively charged ion (e.g., Ca<sup>++</sup>) held on cation exchange capacity (CEC) or in soil solution. Chelate = metal ion held by an organic compound through multiple bonds. Mineral = present in insoluble mineral or mineral-fixed forms. Organic = integral part of soil organic matter (e.g., N and S in proteins or amino acids).

While all nutrients are important to soil, plant, animal, and human health, the remainder of this webinar will focus on nitrogen and phosphorus, which have the greatest potential impact on water resources, soil health, and greenhouse gas emissions.

*Slide 10 – 21<sup>st</sup> Century nutrient management: feed the plant and the soil.*

By the end of the 20<sup>th</sup> Century, most land grant university and other mainstream agricultural professionals recognized the importance of beneficial soil organisms, soil organic matter (SOM) and soil health to long term sustainability of agricultural production and considered biologically derived as well as soluble nutrient sources in making nutrient management recommendations. The role of soil life shifted from mere competitor for nutrients to mediator of a two-way process of immobilization (which can protect water quality and conserve the nutrients for future use as well as reducing short term availability) and mineralization (releasing soluble, crop-available nutrients).

Standard nutrient recommendations now credit manure applications, cover crops, crop residues, and sometimes estimates of N mineralization from SOM. Application of the NRCS “4Rs” leads to more conservative application rates, targeted timing and placement, and a wider range of fast and slow-release nutrient sources, thus reducing losses and environmental harms.

Diversified crop rotations help nutrient management through complementary nutrient demands (root depth, spread, and architecture) and greater microbial functional diversity (legumes, grasses, crucifers, other forbs supporting different root microbiomes). Enhancing biodiversity is one of the four NRCS principles of soil health management, yet a large percentage of US cropland remains in low-diversity rotations like corn-soy or wheat-fallow.

*Slide 11 – In organic farming systems, soil organisms drive nutrient cycling and crop nutrition*

Research over the past 20 years has shown the central role that the soil microbiome plays in all aspects of soil function, and especially nutrient cycling and crop nutrition (red highlighted items in the slide). Other soil health functions – water storage, structure, disease suppression – support crop nutrition, as roots require adequate moisture and oxygen to absorb nutrients, and roots damaged by pest nematodes or microbial pathogens cannot absorb nutrients effectively.

*Slide 12 – Organic nutrient management*

Organic farmers rely primarily on biological processes to meet crop nutrient needs, and supplement with NOP-allowed mineral and faster-releasing organic fertilizers. These more concentrated nutrient sources may be essential during the first few years under organic management, after which the need diminishes as soil health improves.

As the soil life converts plant residues, manure, and other organic inputs into active and stable organic matter, most of the N, P, and S in the residues become integral parts of the organic matter and are slowly released to plants through further action of soil organisms on the active fraction. K, Ca, Mg, and some micronutrients are released from residues into the soil as soluble cations. Negative charges on soil clays and stable, mineral-associated organic matter (MAOM) – the soil’s cation exchange capacity (CEC) – adsorb and hold the cations in a plant-available yet not readily leachable form. In converting some of the residues and active SOM into MAOM, soil organisms maintain and enhance the CEC, an important aspect of soil fertility.

In addition, soil minerals hold large nutrient reserves, particularly potassium (K), other cations, and micronutrients, which are gradually brought into the exchangeable (plant available)

pools through the action of soil life and plant roots on the mineral component of the soil (biological weathering).

The capacity of the soil life to provide for crop nutrition through these processes is a key attribute of healthy agricultural soils. One notable aspect of soil health and plant nutrition is the depth of plant root accessible soil profile. While biological activity is slower at depths below 6–12 inches, plant roots and their associated microbiomes can grow as deep as five feet or more, retrieving leached nutrients (N, S, sometimes others), accessing K and other nutrients from soil mineral reserves, and forming long-lived subsoil MAOM.

Research has shown that the organic method is not “immune” to unwanted nutrient losses including runoff, leaching, and N<sub>2</sub>O emissions; however, studies are identifying promising strategies for maximizing nutrient efficiency and minimizing losses and nutrient pollution.

Most organic farms import some organic amendments from off-farm sources, including manure, compost, municipal leaves, yard trimmings, and food scraps, to replenish nutrients and organic matter consumed in production or removed in harvest.

#### Slide 13 – *Nutrients for carbon: an ancient partnership*

These biological processes can sustain plant nutrition even when soil tests show suboptimal (low to medium) levels of soluble plant-available nitrogen (PAN), phosphorus (P) and other nutrients in the bulk soil. This phenomenon of *tightly coupled nutrient cycling* can protect water quality and mitigate greenhouse gas emissions. Recent paleontological findings indicate that the first land plants co-evolved with mycorrhizal symbionts that proved essential to their survival on the prehistoric world’s primitive soils.

#### Slides 14-16 - *Soil microbes need a “balanced diet” to do their job: the critical role of the carbon-to-nitrogen (C:N) ratio.*

Soil microbes thrive and become most efficient in converting residues into new microbial biomass and active and stable SOM when the residues have a weight ratio of carbon to nitrogen (C:N ratio) between 25:1 and 35:1. When the soil is fed residues with a higher C:N ratio, microbial growth is N-limited and likely constrained by scarcity of other nutrients as well. Since they must “burn off” the excess carbon, a higher proportion of the organic material is converted back into CO<sub>2</sub>, and SOM builds only slowly (Grandy and Kallenbach, 2015).

Most important for the organic farmer to consider is that plants take up nitrogen mainly in the soluble nitrate and ammonium forms, thus “plant available nitrogen” or PAN = nitrate-N + ammonium N. Crops must have access to PAN, either directly from soluble fertilizers, or via mineralization of organic N in active soil organic matter, manure, cover crops, and other organic materials. Mineralization is a process mediated by the soil life, which thus acts as a “gate-keeper” in nitrogen cycling in organic systems. When N-poor residues are incorporated into the soil, microbes must reverse this process, utilizing soluble soil N to make up the deficit, immobilizing it in organic matter and microbial biomass and leaving less available for crop growth. Organic N fertilizer will be needed to avoid crop N deficiency and sustain yields.

When high C:N residues are left on the soil surface (e.g., straw or chipped brush mulch or roll-crimped cereal cover), N immobilization is less intense, and the mulch can benefit soil and crops by conserving moisture, protecting the soil surface from crusting and soil organisms from direct sun and temperature extremes, and suppressing weed seeding emergence. Broccoli transplanted no-till into mowed cereal rye cover (photo) has suffered N deficiency because the rye took up much of the plant available soil N.

When manure, succulent plant residues, or organic fertilizers with a C:N ratio below 15:1 are incorporated into the soil, a rapid release of PAN takes place, which can facilitate high yields in the next crop – with potential environmental costs. Because the soil life is carbon-limited, it must consume some of the carbon in active SOM in order to grow and reproduce, so that a net loss of soil carbon can occur. Furthermore, if nitrate-N is released faster than crops can utilize it, heavy rain or irrigation can leach it below the reach of crop roots so that it is lost from the production system and may pollute groundwater. In addition, soluble soil N is subject to microbial denitrification whenever soil moisture is high and aeration is limited. Denitrified N enters the atmosphere as elemental N<sub>2</sub> gas (harmless other than the wasted fertilizer N) and N<sub>2</sub>O, a powerful greenhouse gas.

In well managed organic systems, the “gatekeeper” – the soil food web – can moderate though not eliminate these losses and environmental impacts. When a high biomass all-legume cover crop is tilled in, or when a long history of heavy organic inputs has built a *very* large pool of active organic matter, N losses can be as great as in high-input conventional systems.

The challenge is to ensure that adequate PAN is available in the crop root zone, and at the same time minimize nitrate leaching and denitrification.

When soil microbes consume organic materials with a balanced C:N ratio, they become more efficient in generating new microbial biomass and active and stable SOM, and mineralize N at a slow, steady trickle that crops are more likely to utilize, leaving less to be lost to leaching, runoff, or denitrification. Using a variety of organic inputs with varying C:N ratios and a composite (average) ratio in the 25 – 35:1 range may be especially beneficial to multiple soil functions including nutrient cycling and retention (Bhowmik et al., 2016, 2017).

Finished compost consists of mixed organic residues that have already been processed by microbes, often in a thermophilic process (130-150°F) that rapidly converts the residues into a mixture of active and stabilized SOM. In a well-managed composting system, about half of the original carbon is converted to CO<sub>2</sub> while most of the N and other nutrients are retained and stabilized, resulting in a lower C:N ratio, perhaps 15:1 or 20:1. Unlike raw organic residues with this low a ratio, finished compost will mineralize only about 10 – 25% of its N to the current year’s crop; however its content of active and stable SOM, other nutrients including P, cations, and micronutrients, and beneficial microbes will enhance soil fertility.

Note that a “balanced diet” for microbes includes all the nutrients discussed earlier – not just C, N, and P. For example, N-fixing and N cycling microbes require several micronutrients including Fe, Mo, Mn, and Cu. Plant residues, manure, compost, and organic fertilizers provide most of these nutrients in varying amounts, while soluble fertilizers essentially consist of “pure” NPK and can leave crops deficient in S, Mg, or micronutrients.

#### Slide 17 – *The whole is greater than the sum of the parts*

One of the hallmarks of the organic method is *systems thinking*. Rather than relying on a single input or practice, successful organic farmers combine diverse crop rotations, cover crops, compost and/or other organic inputs, and judicious tillage to build the soil’s capacity to nourish crops, thereby reducing the need for costly organic fertilizers. Research has shown that “stacking practices” builds more SOM and better soil functions than single practices (Delate et al., 2015; Hurisso et al., 2016).

Organic systems emphasize agricultural diversity, including long rotations that include multiple plant families, intercropping and relay planting, and integrating livestock into cropping systems to support a more diverse and complete soil food web and make best use of soil nutrients. Recent studies have found that intercropping legumes with vegetable crops can modify the soil microbiome to enhance N cycling and reduce fertilizer needs (Cuartero et al., 2022; Stratton et al., 2022).

Organic no-till, even rotational no-till, can be challenging because of yield tradeoffs related to N limitation and weed pressure. However, a recent meta-analysis has shown that reduced tillage – shallow, non-inversion tillage such as the high-speed disk working at 3 – 4 inches, can maintain twice the microbial biomass as either moldboard plowing or continuous no-till with chemical weed control and inadequate cover cropping (Morugán-Coronado et al., 2022).

## Organic Nutrient Management Challenges (Slides 18-45)

Research findings and applying the 4Rs to organic systems

### Slide 19 – *Organic advantages and challenges*

Organic rotational no-till systems that integrate high-biomass cover crops terminated by roller-crimping, diverse crop rotation, and organic soil amendments can maximize soil health parameters such as total and active SOM, soil organic N, and microbial biomass and diversity. However, yield tradeoffs can be severe, especially in colder or drier climates, with up to 63% yield losses in corn and oats related to inadequate PAN and weed competition (Carr et al., 2020; Delate, 2013). However, as noted above, shallow noninversion tillage integrated with cover crops and sound organic soil management can maintain healthy, biologically active soils.

Farmers undertaking organic transition in fields with a history of reliance on soluble fertilizers with inadequate organic inputs face additional challenges in providing for crop nutrition. Such soils often have depleted soil microbiomes and limited active SOM pools, and therefore cannot meet crop N needs from cover crops, compost, and SOM mineralization until soil health is restored. Strategies for the transition period include:

- Increase use of concentrated organic fertilizers such as poultry litter to sustain yields. This approach can delay restoration of SOM and soil health, deter development of microbial N mineralization potential, and perpetuate reliance on expensive inputs.
- Begin with less nutrient-demanding production crops, rotate with cover crops, use compost to restore active SOM and concentrated organic fertilizers in moderation.
- Focus on soil restoration with continuous cover or sod crops + organic carbon amendments (compost, biochar, etc) for a few years before attempting organic crop production. This will rebuild soil health and soil microbiomes most quickly, but not all farms can afford to forego income for this long.
- Integrate livestock into the system to obtain grazing value from cover or sod crops. Sound rotational grazing can facilitate soil restoration.

Relying on manure and compost to meet crop N needs can build up surplus soil P. The complexities of biologically mediated soil nutrient dynamics can create challenges in determining how much N to apply in organic production and synchronizing the timing of N mineralization with crop N need. These challenges will be explored in the following slides.

Slides 20-28 – *The first R: right source*

Slide 21 – *Organic versus soluble – what does the research show?*

The goal of this literature review slide is to address the question “why organic?” through a nutrient management lens.

While soluble NPK fertilizers have boosted crop yields and thereby potential organic matter return to the soil via root exudates and residues, a University of Illinois review of the Morrow Plots and 25 other farming systems trials around the world revealed a downward trend in total SOM and soil organic nitrogen reserves (Khan et al, 2007; Mulvaney et al, 2009).

In a worldwide review of multiple meta-analyses comparing the impacts of different production systems and inputs on soil carbon and nitrogen dynamics, yields, and environmental impacts, Young et al (2021) found that, compared to soluble N, the use of organic sources of N substantially enhances total SOM, curbs N leaching and ammonia (NH<sub>3</sub>) volatilization, but slightly reduces yields and may increase N<sub>2</sub>O emissions. Managing biologically active for N<sub>2</sub>O mitigation is a significant challenge and will be explored in depth in a future webinar on organic systems and climate change.

Compared to soluble fertilizers, the use of organic nutrient sources roughly doubles soil microbial biomass with substantial increases in both fungal and bacterial communities (Morugán-Coronado et al., 2021), and also doubles the biomass of bacterial- and fungal-feeding nematodes (Puissant et al., 2021). These nematodes play a vital role in N mineralization and crop N nutrition in organic systems.

Studies at multiple sites have shown that organically managed soils have substantially greater ability to provide crop-available N through microbial mineralization from SOM (Berthong et al., 2013; Franzluebbers et al., 2018b, 2020; Spargo et al., 2011). This capacity is sustained by the replenishment of organic C and N through cover crops and amendments, which suggests that organic nutrient management can be based on total N rather than PAN.

Other studies have documented qualitative differences in the microbial communities under organic vs soluble fertilizer regimes, with greater biodiversity, improved N and P cycling, and reduced disease pressure with organic nutrient sources (Li et al., 2022; Zhang et al., 2022).

Slide 22 – *Three organic nutrient sourcing strategies*

Best nutrient management combines all three strategies – grow fertility in place (cover crops), on-farm cycling. Each one alone has its pitfalls:

- Cover crops can meet the N needs of the following crop in warm, rainy climates, but may not in cold-temperate (Northeast, Great Lakes) or dry regions (Intermountain West), where N mineralization is slower (Carr et al., 2020). Plowing a perennial legume sod can meet the needs of a following corn crop, but N<sub>2</sub>O emissions can occur (Han et al. 2017).
- On-farm cycling can sustain production for years; however, nutrients exported in farm products sold will eventually have to be replenished with off-farm inputs.
- Generous applications of compost and manure can rebuild depleted soils and support intensive organic production; however, P and other nutrient excesses can accrue, and high compost rates are not economically feasible at larger scale.

Our society wastes huge amounts of organic residue in ways that turn a vital resource into an environmental problem: livestock manure and urine (lagoons or stockpiles), yard trimmings, leaves, and food scraps (landfill). All of this can and should be composted and returned to the land to replenish soils and reduce fertilizer needs for crops.

Slide 23 – *Cover crops: a vital organic nutrient management tool.*

Cover crops play vital, multiple roles in soil-friendly nutrient management. As a key source of organic carbon to feed soil life and maintain soil organic matter, high biomass cover crops enhance the soil's long term capacity to provide for crop nutrition. Winter grass and legume cover crops provide a “green bridge” to sustain mycorrhizal populations.

Cover crops help to regulate soil nutrient levels. When soil soluble N is scarce, legumes maximize N fixation, and some warm season grasses host N-fixing bacteria in their root zone. When soil soluble N is abundant, these cover crops switch to “scavenging mode,” absorbing and holding the surplus N.

When plant-available P or K are below optimum, cover crops can enhance their availability. Buckwheat and most legumes can retrieve P from insoluble organic and mineral sources (including rock phosphate amendments), while most grasses can unlock “mineral-fixed” K to replenish the supply of exchangeable (plant-available) K. However, cover crops do not “fix” P and K from thin air the way they do C and N – thus, cover crops will not add unneeded P and K when soil levels are already ample.

Slide 24 - *On-farm Nutrient Sourcing Elmwood Stock Farm, Georgetown, KY*

University of Kentucky scientists have studied the integrated farming system at Elmwood Stock Farm and found that the five-year sod phase under rotational grazing restores SOM and microbial community to close to that of permanent pasture (Lin et al., 2020). However, when the sod is broken to resume crop production, a burst of SOM oxidation and N<sub>2</sub>O emission occurs (Shrestha et al., 2019). Beginning in 2019, the farmers have experimented with grazing and shallow tillage to end the sod phase with less soil disruption, and with alternative crop rotations to optimize production, soil health, and GHG mitigation outcomes.

Slide 25 – *Off-farm organic nutrient sources: which materials are best?*

A research team at Washington State University compared the crop and soil impacts of two nutrient sources in organic vegetable production in a maritime soil in Washington State: on-farm mixed compost made from dairy manure and bedding and yard waste (C:N ~20) at rates of 6 to 8 tons/ac annually, and composted poultry litter (C:N ~7) at 1.8 – 2.6 tons/ac annually. The total N amounts applied in the two treatments were similar. Crop yields, soil physical, chemical, and biological properties, and potential N<sub>2</sub>O emissions were monitored over an 11-year period.

The higher C:N compost improved overall soil health, with substantially higher levels of active and total SOM, microbial activity, and enzyme activities involved in nutrient cycling; and a more balanced nematode community (Bhowmik et al., 2016, 2017; Cogger et al., 2013). Notably, the compost amended soil showed both a greater capacity to mineralize N for crop production and to immobilize excess soluble N. This suggests that reliance on concentrated, low C:N inputs like poultry litter may shift the soil microbiome in a way that weakens N cycling and provisioning, thereby perpetuating reliance on these inputs.

Slide 26 – *Nutrient source and NPK balance: vegetable crops*

Typical yields, and estimates for N (lower figures), P, and K removal based on nutrient concentrations given in Knotts Handbook for Vegetable Growers, 5th ed (D. N. Maynard and G. J. Hochmuth, 2007) cited in Wander (2009).

The higher values for N removals are based on vegetable nutritional contents (N = protein / 6.25) cited in Nutrition Almanac, 3rd edition (Donne, 1990).

Extension recommendations for N (regardless of soil test) and for P and K at “high” (optimum) soil test P and K levels are from Virginia Tech (2018).

Nutrient replenishment to maintain optimum soil nutrient levels should aim to balance inputs with exports through harvest. NPK recommendations for fertility maintenance seem appropriate for K, high for N (especially head brassicas), and severalfold too high for P.

Note the low nutrient removal rates for vegetable harvests, especially P. Moderate rates of compost and poultry litter – two amendments commonly used for NPK in organic vegetable production – can approximate Extension recommendations but will build up soil P over time.

#### *Slide 27 – Nutrient source and NPK balance: field crops*

Nutrient removal estimates for the yields shown are based on Virginia Tech Extension data for nutrient contents per bushel of grain or ton of forage.

Grain crops remove relatively more N and P, and less K, compared to vegetables. Most of the K taken up by the crop is returned in residues. Forage harvests remove substantial amounts of all three nutrients.

Compost and poultry litter applied at rates sufficient to replenish N will build up P and (for grains) K.

#### *Slide 28 – Mix and match sources to get the balance right*

Rotating vegetables, grains, and forages with complementary nutrient demands can help balance soil levels and facilitate nutrient budgeting. Growing vegetables with compost or poultry litter year after year will build excess P, while rotating to hay can help draw down the P surplus. Haying year after year can deplete soil P and K reserves, whereas grazing recycles half or more of the nutrients via manure/urine deposition.

#### *Slides 29-33 – The second R: right rate*

##### *Slide 30 – Total versus Available N*

Rates for organic nutrient resources are often based on their “available” N percentage, the fraction of N in the amendment that moves into the crop during the season of application. However, this can lead to costly and potentially polluting over-application and GHG emissions (Baas et al., 2015). Because soil biological activity mineralizes N and other nutrients from active SOM to support crop growth, it may only be necessary to replenish the SOM reserves to balance N removals in harvest, and therefore to use organic N sources at rates based on their total N content. A recent global meta-analysis of 129 studies (Wei et al., 2021) indicates that this may be true. In comparisons of organic versus conventional soluble N sources, organic applications based on *total* N maintained yields and reduced N leaching and runoff losses by an average of 30%. Organic applications based on *soluble* N content nudged yields an average of 6% above conventional yields, but also resulted in N losses 21% higher than the conventional treatment.

##### *Slide 31 – Grain crops may need little fertilizer on healthy soils*

Fertilizer trials were conducted over a five-year period with an organic corn-soy-wheat rotation with legume-cereal cover crops on an Orangeburg loamy sand in South Carolina (Kloot, 2017, 2018). The cover crops attained high biomass and substantial nutrient accumulations: 9000

lb dry matter, 110 lb N, 27 lb P, and 200 lbK per acre, which allowed the crops to attain full yields on half-rate N and no P or K. Furthermore, soil test P and K levels showed little or no decline during the five years, and Kloot noted that many soils have tremendous subsoil K reserves that grass cover crop roots can access. Similarly, farmers in a diversity of locations (NC, ND, IL, OH) have greatly reduced fertilizer inputs and maintained high grain yields by building soil health with high biomass cover crops.

Slide 32 – *Economic optimum nitrogen rate (EONR) for crops in healthy soils can drop to zero*

Researchers at North Carolina State University have documented a close correlation between “soil test biological activity” (STBA, a three day soil microbial respiration test) and the soils’ capacity to make N available to crops through SOM mineralization, and conducted N fertilizer-yield response trials at multiple locations. Farmer fields with high STBA, managed organically or in accord with the NRCS soil health principles, showed so little response to added N that their EONR was zero for corn grain, silage, or fescue forage production (Franzluebbers, 2018; Franzluebbers et al., 2018a). Similarly, high biomass rye-clover cover crops (total N 130 b/ac) supported good organic tomato and summer squash yields with no response to added N, even when the vegetables were no-till planted into roll-crimped covers (Robb & Zehnder, 2016).

In a review of multiple field trials, Khan et al (2013) found that standard K recommendations are far higher than necessary, and sometimes lead to decreased soil health or crop quality. Based on this finding, organic producers with soils testing below optimum in K may not need nearly as much potassium sulfate ( $K_2SO_4$ ) as suggested by standard soil test recommendations. Using less will save growers money and protect soils from salinity burdens.

Slide 33 – *EONR soars above 200 lb/ac for organic broccoli*

Field trials at University of California Santa Cruz evaluating organic broccoli yield response to 0, 75, 150, and 225 lb N/ac in the form of blood, meat, and feather meals, with or without compost or legume-cereal cover crop, indicated an economic optimum nitrogen rate of 215 lb/ac (Li et al., 2009). The trial also included studies of soil C and N dynamics, which led to estimates of major N losses.

Trials on five organic farms in coastal and interior Oregon (seven site-years) confirmed that organic broccoli yields show a significant linear response to added N (as feather meal) of 11 to 88 lb marketable yield per lb N, and did not plateau until  $> 200$  lb N/ac (Collins and Bary, 2017). At a market price of \$2.50/lb, broccoli yield increases returned several times the cost of the feather meal (about \$6/lb) at all sites. Thus, the high market value of organic broccoli, combined with the strong yield responses resulted in a high EONR.

Why was organic broccoli so inefficient in N uptake, leaving so much in the field to leach? Broadcast application of fertilizers on a crop with a fairly narrow lateral root spread may have been a factor. In addition, the Mediterranean climate pattern of winter rain and dry summer (broccoli grown during the dry season with irrigation) may have slowed mineralization from the organic fertilizers, then promoted leaching in winter without cover crops. However, other trials by Virginia Tech in Blacksburg, VA (rainy summer) showed yield responses to at least 150 lb N/ac in addition to the N from a roll-crimped vetch + rye cover crop.

Slides 34-39 – *The third R: right timing*

Precision timing through split applications is straightforward for soluble N sources and challenging for organic sources that must undergo biological processing before plant uptake.

Slide 35-37 – *Synchronizing N release with crop N demand*

N demand in most annual crops goes through three distinct phases; a “lag” period during the first 3-4 weeks after planting when crop N needs are relatively small, a vegetative phase of rapid plant development and high N demand, and a maturation phase, during which N uptake slows as N is translocated from leaves to developing fruit, grain, or tuber.

N use efficiency is maximized when the release of PAN from fertilizers, amendments, and soil organic matter is synchronized with the period of high N demand, so that crops are not N limited, yet excess soluble N does not remain in the soil for extended periods of time. If N is released too quickly, early season rains may leach it out before the crop can utilize it. N released too slowly becomes available only after the crop has entered the maturation phase, resulting in crop N limitation and an increased risk of late season leaching or denitrification losses.

An organic fertility program that provides ideal crop N nutrition in warm, moist, aerobic, biologically active soil may release N too slowly in cold, wet, or hot dry conditions, or if soil life and soil health are below par.

For example, organic no-till corn planting into roll-cripped cover crops often fails in the upper Midwest. Even though the region’s Mollisol soils have high inherent fertility, the cool climate, short growing season, and heavy soil textures can delay biological N mineralization so that the corn may not obtain sufficient N from a roll-cripped rye + vetch cover crop, whereas tilling the cover crop gives timely release of PAN and good corn yields.

Conversely, in the southeastern US coastal plain, warm rainy climate, long growing season, and sandy Ultisol soils promote rapid N mineralization – possibly too rapid if the cover crop is tilled in. In these conditions, a roller-cripped rye + vetch cover may provide more timely N delivery to the crop.

Slide 38 - *Asynchrony of N supply and N demand in an organic strawberry field in the Northern region, CA*

This slide is taken from a 2015 webinar with permission from Dr. Joji Muramoto of University of California at Santa Cruz.

Strawberry is a major crop in California, accounting for 85% of US production. 9% of CA strawberry acreage is organic. Strawberries are grown in coastal regions of central CA, where mild, dry summers allow prolonged harvests and high yields.

N management is especially challenging because of the very long lag phase in the crop’s N utilization, which often renders preplant applications of organic N sources ineffective. The strawberry crop in this diagram, planted in November after broccoli harvest, was unable to utilize some 260 lb nitrate-N mineralized from broccoli residues and pre-plant organic fertilizer. Winter rains leached most of this N out of the root zone long before the strawberry crop began taking up N more rapidly in late spring (Gaskell et al., 2009; Muramoto et al., 2015). Preplant N from other sources such as summer cover crops and compost is similarly mineralized and lost during the winter.

Slide 39 – *Organic vegetables + cover crop*

While strawberry planted after broccoli cannot utilize the leftover N, vigorous, deeper-rooted winter cover crops can do so quite effectively.

Dr. Eric Brennan of USDA Agricultural Research Service in the Salinas Valley of California conducted an eight-year trial (Salinas Organic Cropping Systems Experiment) on a Chualar

loamy sand, which is well drained but has a root-restrictive layer at about 30 inches below the surface. A double cropping system of spring lettuce followed by fall broccoli sustained high lettuce yields (1000 boxes/ac, about 30 lb/box) only when a winter cover crop was grown prior to the lettuce. Whenever the field was left fallow over winter, lettuce yields declined sharply to a few hundred boxes per acre, and sometimes to a total crop failure. Cover crops of rye alone, mustard, or rye with vetch, fava, and pea were similarly effective, indicating that their main benefit was not N fixation per se, but recovery of N left over from the broccoli crop. Broccoli was fertilized with about 145 lb N/ac (from NOP allowed organic sources), only about 25% of which was removed in harvest. During winter fallow, winter rains leached the N from the entire root-accessible zone, whereas vigorous winter cover crops recovered N and their residues delivered it to the lettuce (Brennan, 2018).

In addition, the combination of cover crops plus compost enhanced soil microbial activity to a much greater degree than the compost alone, even though the compost comprised the majority of organic carbon inputs (Brennan & Acosta-Martinez, 2017).

#### Slides 40-43 – *the fourth R: right placement*

Band application or in-row drip fertigation can deliver N and other nutrients right in the root zone. In biologically active soils, microbes may help with right placement of soluble N.

#### Slide 41 – *Delivering N directly to the roots*

What if the “gatekeeper” of soil life could deliver soluble N directly to plant roots as they need it, without flooding the bulk soil with soluble N that would be subject to leaching and denitrification? This happens in the legume-Rhizobium symbiosis, in which atmospheric elemental nitrogen is “fixed” into plant-available N to the benefit of both symbionts, as shown by the soybean in this illustration. Non-leguminous plants can also host N-fixing bacteria, though not in visible nodules. For example, landraces of corn have been identified that derive up to 40% of their N requirement from N fixing rhizosphere microbes, and plant breeders have successfully transferred this trait to modern grain corn hybrids, enabling them to thrive and yield in low-N soils and on greatly reduced fertilizer rates (Goldstein, 2015, 2016, 2018). Some other warm-season grasses including pearl millet can also host N-fixing microbes to meet part of their N requirement.

Can other soil organisms help plants access the large store of soil organic N by mineralizing just the right amount at the right time in their root zone? There is strong evidence that this can happen. Rhizosphere (root zone) microbial population densities are typically 10 times those of bulk soil. Plant roots give off chemical signals that attract the “right” microbes into their vicinity, then feed them with nourishing root exudates. In turn, the microbes help the plants acquire the moisture, N, and other nutrients they need, increase plant resilience to stress and diseases.

#### Slide 42 – *N cycling patterns in organic tomato in California*

The slide summarizes findings from a study of 13 organic tomato fields in central California.

The four fields that showed “tightly coupled nitrogen cycling” with low soil soluble N levels yet adequate crop nutrition and high yields were amended primarily with a fairly high C:N compost (20:1) derived from diverse organic materials mixed together. This was supplemented with a light in-row application (band or drip fertigation) of more concentrated N (blood meal or Chilean sodium nitrate). The soils had also been under organic management with cover crops for a number of years. The investigators documented evidence that the soil microbiome and plant

root enzymes interacted to enhance N availability within the rhizosphere without overloading the bulk soil with soluble N (Bowles et al., 2015; Jackson, 2013; Jackson & Bowles, 2013).

In contrast, two N-deficient fields with less healthy soils were unable to mineralize sufficient N for crop nutrition, while seven N-saturated fields that received most of their N in relatively concentrated (low C:N) forms showed high microbial activity but with lower levels of N cycling enzymes and higher levels of enzymes involved in SOM oxidation.

The authors stated that, since the plant root enzymes involved in tightly coupled N cycling occur widely across crop species, this phenomenon could occur in many crops and locations.

#### Slide 43 – *How to promote root zone N mineralization*

Beneficial microbes thrive in the rhizosphere when plants deliver sufficient root exudates. In a research review, Prescott et al (2021) identified three strategies to promote root exudation and thereby enhance carbon sequestration in the form of mineral-associated organic matter:

- Keep plant available N and P levels, and irrigation levels slightly below the optimum for top growth. These slight deficits do not inhibit photosynthesis, and therefore they create a surplus of photosynthetic product that the crop sends into the root zone and the rhizosphere as root exudates. Yields are not adversely affected, and may benefit from enhanced root growth and rhizosphere microbiome.
- Include legumes in the crop rotation or pasture mix. Unlike soluble N fertilizers, the N-rich root exudates from legumes provide excellent nutrition for soil microbes.
- Time rotational grazing events late in the rapid growth phase of the forages, as the rapid growth phase is also the time of greatest root exudation.

The authors did not evaluate impacts on N cycling; however, providing sufficient microbial “food” in the form of root exudates seems likely to enhance this microbial function as well.

Avoiding N and P excesses and limiting use of concentrated (low C:N) organic N sources appears important for sustaining the desired soil microbiome. However, the California tomato study suggests that, if crops need a boost (e.g., in early development before they develop a deep and extensive root system), using small doses of concentrated N in the row may be compatible with efficient biological N cycling.

Finally, there is growing evidence that crop cultivars have genetic differences in their capacity to partner with beneficial microbes, and that crops can be bred and selected for nutrient efficiency. The USDA Organic Research and Extension Initiative is funding several farmer-participatory plant breeding endeavors that are exploring this potential.

#### Slide 44 – *Summary of organic nutrient management tips*

## References

Baas, D. G., G. P. Robertson, S. R. Miller, N. and Millar, N. 2015. Effects of Cover Crops on Nitrous Oxide Emissions, Nitrogen Availability, and Carbon Accumulation in Organic versus Conventionally Managed Systems. Final report for ORG project 2011-04952.  
<https://nifa.usda.gov/data/data-gateway>.

Barbercheck, M. E. 2016. A Reduced-Tillage Toolbox: Alternative Approaches for Integrating Cover Crops and Reduced Tillage in an Organic Feed and Forage System. Project proposal and progress report for OREI project 2014-05377. <https://nifa.usda.gov/data/data-gateway>.

Barbercheck, M. E., W. Curran, J. Harper, R. Hoover, D. Voight, and G. Hostetter. 2014. Improving Weed and Insect Management in Organic Reduced-Tillage Cropping Systems. Final report on OREI project 2010-03391. <https://nifa.usda.gov/data/data-gateway>.

Berthong, S. T, D. H. Buckley, and L. E. Drinkwater. 2013. Agricultural management and labile carbon additions affect soil microbial community structure and interact with carbon and nitrogen cycling. *Microbial Ecology* 66: 158-170.

Bhowmik, A. A-M. Fortuna, L. J. Cihacek, A. Bary, P. M. Carr, and C. G. Cogger. 2017. *Potential carbon sequestration and nitrogen cycling in long-term organic management systems*. *Renewable Agriculture and Food Systems*, 32 (6): 498-510.

Bhowmik, A., A. Fortuna, L. J. Cihacek, A. I. Bary, and C. G. Cogger. 2016. *Use of biological indicators of soil health to estimate reactive nitrogen dynamics in long-term organic vegetable and pasture systems*. *Soil Biology and Biochemistry* 103: 308-319.

Bowles, T. M., A. D. Hollander, K. Steenwerth, and L. E. Jackson. 2015. *Tightly-Coupled Plant-Soil Nitrogen Cycling: Comparison of Organic Farms across an Agricultural Landscape*. PLOS ONE peer-reviewed research article. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0131881>

Brennan, E. 2018. Lessons from long-term, cover crop research in the Salad Bowl of the World – 10 minute youtube video, <https://www.youtube.com/watch?v=JurC4pJ7Lb4>

Brennan, E. B., and V. Acosta-Martinez. 2017. Cover cropping frequency is the main driver of soil microbial changes during six years of organic vegetable production. *Soil Biology and Biochemistry* 109: 188-204.

Carr, P. M, M. A. Cavigelli, H. Darby, K. Delate, J. O. Eberly, H. K. Fryer, G. G. Gramig, J. R. Heckman, E. B. Mallory, J. R. Reeve, E. M. Silva, D. H. Suchoff, and A. . Woodley. 2020. Green and animal manure use in organic field crop systems. Review article. *Agronomy Journal* 112 (2): 648-674.

Clark, K. 2016. Organic weed management systems for Missouri. Proposal and progress report on OREI project 2014-05341. <https://nifa.usda.gov/data/data-gateway>.

Cogger, C. G. M. Ostrom, K. Painter, A. Kennedy, A. Fortuna, R. Alldredge, A.; Bary, T. Miller, D. Collins, J. Goldberger, A. Antonelli, and B. Cha. 2013. Designing Production Strategies for Stewardship and Profits On Fresh Market Organic Farms. Final report for OREI project 2008-01247. <https://nifa.usda.gov/data/data-gateway>.

Collins, D. P. and A. Bary. 2017. *Optimizing nitrogen management on organic and biologically intensive farms*. Proceedings of the Special Symposium on Organic Agriculture Soil Health Research at the Tri-Societies Annual Meeting, Tampa, FL, October 22-25, 2017. <http://articles.extension.org/pages/74555/live-broadcast:-organic-soil-health-research-special-session-at-the-tri-societies-conference>.

Cuartero, J., J. A. Pascual, J-M Vivo, O. Özbolat, V. Sánchez-Navarro, M. Egea-Cortines, R. Zornoza, M. M. Mena, E. Garcia, and M. Ros. 2022. A first-year melon/cowpea intercropping system improves soil nutrients and changes the soil microbial community. *Agriculture, Ecosystems, and Environment* vol. 328 article 107856.

Delate, K. 2013. Developing Carbon-positive Organic Systems through Reduced Tillage and Cover Crop Intensive Crop Rotation Schemes. Final report for ORG project 2008-01284. <https://nifa.usda.gov/data/data-gateway>.

Delate, K., C. Cambardella, and C. Chase. 2015. *Effects of cover crops, soil amendments, and reduced tillage on carbon sequestration and soil health in a long-term vegetable system*. Final report for ORG project 2010-03956. <https://nifa.usda.gov/data/data-gateway>.

Donne, L. J. 1990. *Nutrition Almanac*, 3<sup>rd</sup> Edition. McGraw-Hill, 339 pp.

Franzluebbers, A. J. 2018. *Soil-Test Biological Activity with the Flush of CO<sub>2</sub>: III. Corn Yield Responses to Applied Nitrogen*. *Soil Science Society of America Journal*, Volume 82, Issue 3, <https://doi.org/10.2136/sssaj2018.01.0029>.

Franzluebbers, A. J., S. Pehim-Limbu, and M. H. Poore. 2018a. *Soil-Test Biological Activity with the Flush of CO<sub>2</sub>: IV. Fall-Stockpiled Tall Fescue Yield Response to Applied Nitrogen*. *Agronomy Journal*, Volume 110, Issue 5, <https://doi.org/10.2134/agronj2018.03.0146>.

Franzluebbers, A. J., M. R. Pershing, C. Crozier, D. Osmond, and M. Schroeder-Moreno. 2018b. *Soil-Test Biological Activity with the Flush of CO<sub>2</sub>: I. C and N Characteristics of Soils in Corn Production*. *Soil Science Society of America Journal*, Volume 82, Issue 3, <https://doi.org/10.2136/sssaj2017.12.0433>.

Franzluebbers, A. J., S. C. Reberg-Horton, and N. G. Creamer. 2020. *Soil carbon and nitrogen fractions after 19 years of farming systems research in the Coastal Plain of North Carolina*. *Soil Science Society of America Journal*, Volume 84, pp 856-876.

Gaskell, M., M. Bolda, J. Muramoto, and O. Daugovish, 2009. *Strawberry Nitrogen Fertilization from Organic Nutrient Sources*. *Acta Horticulturae* (ISHS) 842:385-388.

Goldstein, W. 2015. Breeding corn for organic farmers with improved N efficiency/N fixation, and protein quality. Proceedings of the Organic Agriculture Research Symposium. <https://eorganic.info/node/12972>.

Goldstein, W. 2016. Partnerships between Maize and Bacteria for Nitrogen Efficiency and Nitrogen Fixation. Bulletin 1. Mandaamin Institute, Elkhorn, Wisconsin, 49 pp.  
<http://www.mandaamin.org/about-nitrogenfixing-corn>.

Goldstein, W. 2018. High Methionine, N Efficient Field Corn from the Mandarin Institute/ Nokomis Gold Seed Co. Proceedings of the 9th Organic Seed Growers Conference, Feb 14-17, 2018, Corvallis OR, pp 25-26. <https://seedalliance.org/all-publications/>.

Grandy, S., and C. Kallenbach. 2015. *Microbes drive soil organic matter accumulation in organic cropping systems*. Recording from the Organic Agriculture Research Symposium, LaCrosse, WI February 25-26, 2015, <http://eorganic.info/node/12972>.

Han, Z., M. T. Walter, and L. E. Drinkwater. 2017. Impact of cover cropping and landscape positions on nitrous oxide emissions in northeastern U.S. agroecosystems. *Agriculture, Ecosystems and Environment* vol. 245: pp 124-134.

Hurisso, T. T., S. W. Culman, W. R. Horwath, J. Wade, D. Cass, J. W. Beniston, t. M. Bowles, A. S. Grandy, A. J. Franzluebbers, M. E. Schipanski, S. T. Lucas, and C. M. Ugarte. 2016. Comparison of Permanganate-Oxidizable Carbon and Mineralizable Carbon for Assessment of Organic Matter Stabilization and Mineralization. *Soil Sci. Soc. Am. J.* 80 (5): 1352-1364.

Jackson, L. 2013. *Researcher and Farmer Innovation to Increase Nutrient Cycling on Organic Farms*. Proposal and final report for OREI project 2009-01415.  
<https://nifa.usda.gov/data/data-gateway>.

Jackson, L. and T. Bowles. 2013. *Researcher and Farmer Innovation to Increase Nitrogen Cycling on Organic Farms* (Webinar).  
<http://articles.extension.org/pages/67391/researcher-and-farmer-innovation-to-increase-nitrogen-cycling-on-organic-farms-webinar>.

Khan, S. A., R. L. Mulvaney, T. R. Ellsworth, and C. W. Boast. 2007. *The myth of nitrogen fertilization for soil carbon sequestration*. *J. Environ. Qual.* 36:1821–1832.

Khan, S. A., R. L. Mulvaney, and T. R. Ellsworth. 2013. *The potassium paradox: implications for soil fertility, crop production, and human health*. *Renewable Agriculture and Food Systems*: doi:10.1017/S1742170513000318. 25 pp.

Kloot, Robin. 2017. *Rethinking P and K fertility in coastal plain soils*. Presentation at the 2017 Organic Agriculture Research Symposium, Lexington, KY, January 26, 2017.

Kloot, Robin. 2018. *Using adaptive nutrient management to answer “how much fertilizer do you actually need?”* NRCS webinar May 8, 2018. Science and Technology Training Library, <http://www.conservationwebinars.net/listArchivedWebinars>.

Li, C., Salas, W. and Muramoto, J. 2009. *Process Based Models for Optimizing N Management in California Cropping Systems: Application of DNDC Model for nutrient management for*

*organic broccoli production*. Conference proceedings 2009 California Soil and Plant Conference, 92-98. Feb. 2009. <http://ucanr.edu/sites/calasa/files/319.pdf>.

Li, J., Y. Yang, J. Wen, F. Mo, and Y. Liu. 2022. Continuous manure application strengthens the associations between soil microbial function and crop production: Evidence from a 7-year multisite field experiment on the Guanzhong Plain. *Agriculture, Ecosystems, and Environment* vol. 338 article 108082.

Lin, D., R. L. McCulley, J. L. Nelson, K. J. Jacobsen, and D. Zhang. 2020. Time in pasture rotation alters soil microbial community composition and function and increases carbon sequestration potential in a temperate agroecosystem. *Science of the Total Environment* 698, <https://doi.org/10.1016/j.scitotenv.2019.134233>.

Morugán-Coronado, A., P. Pérez-Rodríguez, E. Insolia, D. Soto-Gómez, D. Fernández-Calvino, and R. Zornoza. 2022. *The impact of crop diversification, tillage and fertilization type on soil total microbial, fungal and bacterial abundance: A worldwide meta-analysis of agricultural sites*. *Agriculture, Ecosystems, and Environment* vol. 329 article 107867.

Mulvaney, R. L., S. A. Khan, and T. R. Ellsworth. 2009. *Synthetic nitrogen fertilizers deplete soil nitrogen: a global dilemma for sustainable cereal production*. *J. Environ. Qual.* 38:2295–2314.

Muramoto, J., C. Shennan, and J., M. Gaskell. 2015. *Nitrogen management in organic strawberries: challenges and approaches*. (Webinar) <https://eorganic.org/node/14818>.

Prescott, C. E., Yi. Rui, M. F. Cotrufo, and S. J. Grayston. 2021. *Managing plant surplus carbon to generate soil organic matter in regenerative agriculture*. *J. Soil & Water Conservation* 76(6): 99A-104A.

Puissant, J., C. Villenave, C. Chauvin, C. Plassard, E. Blanchart, and J. Trap. 2021. *Quantification of the global impact of agricultural practices on soil nematodes: A meta-analysis*. *Soil Biology and Biochemistry* Volume 161, October 2021, 108383.

Robb, D. and G. Zehnder. 2016. Weeds, nitrogen, and yield: measuring the effectiveness of an organic no-till system. Final report for Southern SARE project GS13-126. <https://projects.sare.org/project-reports/gs13-126/>.

Shrestha, D., O. Wendroth, and K. L. Jacobsen. 2019. Nitrogen loss and greenhouse gas flux across an intensification gradient in diversified vegetable rotations. *Nutrient Cycling in Agroecosystems*, <https://doi.org/10.1007/s10705-019-10001-8>.

Smallwood, M., D. Collins, and B. Howell. 2015. Soil Health in Organic Farming Systems, 2015 Organicology Conference selected live broadcasts. <https://eorganic.org/node/12903>.

Snyder, L., M. Schonbeck, T. Velez, and B. Tencer. 2022. 2022 National Organic Research Agenda: Outcomes and Recommendations from the 2020 National Organic & Transitioning

Farmer Surveys and Focus Groups. Organic Farming Research Foundation, <https://ofrf.org>, 232 pp.

Spargo, J. T., M. A. Cavigelli, S. B. Mirsky, J. E. Maul, and J. J. Meisinger. 2011. *Mineralizable soil nitrogen and labile soil organic matter in diverse long-term cropping systems*. *Nutr. Cycl. Agroecosyst* (2011) 90:253–266.

Stratton, A. E., J. J. Comin, I. Siddique, D. R. Zak, L. D. Filipini, R. R., Lucas, and J. Blesh. 2022. Assessing cover crop and intercrop performance along a farm management gradient. *Agriculture, Ecosystems, and Environment* vol. 332 article 107925.

USDA National Organic Program. Organic Regulations. <https://www.ams.usda.gov/rules-regulations/organic>.

Virginia Tech. 2018. Mid-Atlantic Commercial Vegetable Production Recommendations. <https://www.soiltest.vt.edu/Files/handbooks.html>.

Wander, M. 2009. Nutrient budgeting basics for organic farming systems. <https://eorganic.org/node/3060>.

Wei, Z, E. Hoffland, M. Zhuang, P. Hellegers, and Z. Cui. 2021. *Organic inputs to reduce nitrogen export via leaching and runoff: A global meta-analysis*. *Agriculture, Ecosystems, and Environment* 291. <https://doi.org/10.1016/j.envpol.2021.118176>.

Young, M. D., G. H. Ros, and W. de Vries. 2022. *Impacts of agronomic measures on crop, soil, and environmental indicators: A review and synthesis of meta-analysis*. *Agriculture, Ecosystems, and Environment* 319. <https://doi.org/10.1016/j.agee.2021.107551>.

Zhang, H., X. Zheng, X. Wang, and W. Xiang. Effect of fertilization regimes on continuous cropping growth constraints in watermelon is associated with abundance of key ecological clusters in the rhizosphere. *Agriculture, Ecosystems, and Environment* vol. 339 article 108135.