



Genome Sequence of a Lethal Vascular Wilt Fungus, *Verticillium nonalfalfae*, a Biological Control Used Against the Invasive *Ailanthus altissima*

 Matt T. Kasson,^a Lindsay R. Kasson,^b Kristen L. Wickert,^a Donald D. Davis,^c  Jason E. Stajich^d

^aDivision of Plant and Soil Sciences, West Virginia University, Morgantown, West Virginia, USA

^bSchool of Medicine, West Virginia University, Morgantown, West Virginia, USA

^cDepartment of Plant Pathology and Environmental Microbiology, The Pennsylvania State University, University Park, Pennsylvania, USA

^dDepartment of Microbiology and Plant Pathology and Institute for Integrative Genome Biology, University of California, Riverside, California, USA

ABSTRACT *Verticillium nonalfalfae*, a cosmopolitan soil-borne phytopathogen, causes vascular wilt in agricultural crops and perennial woody plants. Select strains of *V. nonalfalfae* can cause lethal disease in the invasive tree *Ailanthus altissima* and several have since been utilized as a biological control (biocontrol) against this widespread invader. Here, we report the genome sequence and annotation of *V. nonalfalfae* strain VnAa140/NRRL 66861.

Since 2005, *Verticillium nonalfalfae* has been recovered from symptomatic *Ailanthus altissima* ("tree-of-heaven") across nine forested locations in the eastern United States. (1–4). These disease epicenters represent natural infections in mixed hardwood stands newly invaded by *A. altissima* (5). More recently, *V. nonalfalfae* has been recovered from dying *Ailanthus* trees in Austria, the first confirmed occurrence on this host outside the United States. (6, 7). This disease is characterized by acute wilting and defoliation followed by epicormic sprouting and mortality, typically within a single growing season (1, 2) (Fig. 1). Several studies using various *V. nonalfalfae* strains recovered from *Ailanthus* trees have helped elucidate phylogenetic relationships (2), biocontrol efficacy (2–4, 8), host range (2, 9–11), and transmission (12) in an attempt to understand if and how they differ from strains impacting agronomic crops. Given the ongoing investigations of *V. nonalfalfae* isolates from hops in Europe (13) coupled with the recent discoveries of *V. nonalfalfae* in Austria, we anticipate that genome-wide comparisons among *Ailanthus* trees and hop-origin strains will provide additional insight into host specificity. There also is a renewed interest in utilizing *V. nonalfalfae* as a biocontrol against *Ailanthus altissima* in the United States to help combat a newly established invasive insect, *Lycorma delicatula*, a planthopper that preferentially feeds on this host (14). The generation of genomic resources, including for biocontrol strains, is fundamental to these efforts.

Verticillium nonalfalfae (strain VnAa140/PSU140/NRRL 66861) was isolated from a dying *A. altissima* in Pennsylvania in 2005 (1, 2). Mycelial fragments and conidia from a 1-week-old culture were transferred to potato dextrose broth and incubated for 2 weeks. Genomic DNA was extracted from the mycelium using a Qiagen DNeasy plant mini kit, and ~5 μg of input genomic DNA was used to construct sequencing libraries. Illumina HiSeq 2000 sequencing generated 29.3 M paired sequence reads or 3.4 Gb (100× coverage), and 454 sequencing produced 1.3 M singleton reads totaling 523 Mb (16× coverage) at the Pennsylvania State University Huck Institutes Genomics Core. The Illumina sequence reads were assembled with Velvet (v0.7.61) (kmer, 31; -cov_cutoff auto; -ins_length 300; -min_contig_lgth 100; -exp_cov auto) (15). Newbler (v2.3) was used to quality trim the 454 reads by default parameters and generate a hybrid 31.7-Mb

Citation Kasson MT, Kasson LR, Wickert KL, Davis DD, Stajich JE. 2019. Genome sequence of a lethal vascular wilt fungus, *Verticillium nonalfalfae*, a biological control used against the invasive *Ailanthus altissima*. *Microbiol Resour Announc* 8:e01619-18. <https://doi.org/10.1128/MRA.01619-18>.

Editor Antonis Rokas, Vanderbilt University

Copyright © 2019 Kasson et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jason E. Stajich, jason.stajich@ucr.edu.

Received 28 November 2018

Accepted 15 December 2018

Published 24 January 2019




FIG 1 Wilt symptoms in *Ailanthus* trees following artificial inoculation with *Verticillium nonalfalfae* strain VnAa140/NRRL 66861, including, stand dieback and mortality (A); acute wilt and defoliation (B); wilted epicormic sprouts that emerged following dieback of the main stem (C); and conspicuous yellow vascular discoloration and streaking (D).

assembly ($n = 630$; N_{50} , 172 kb; max contig, 678 kb; G+C content, 55%) from the Velvet contigs and 454 reads (16). This assembly was cleaned of vector contamination and redundant contigs by Automatic Assembly for the Fungi (AAFTF; v0.2.1) (17) and was further corrected by 5 rounds of polishing with Pilon (v1.22) using the Illumina reads. Genome annotation was performed with funannotate (v1.5.0-760de7c) (18) utilizing available *Verticillium dahliae* and *V. alfalfae* transcripts and proteins as evidence (13, 19). The *ab initio* gene predictor GeneMark.hmm ES was self-trained using the default protocol and Augustus parameters trained from alignments of the BUSCO protein set sordariomyceta_odb9 (20–22), and parameters were archived in a public repository (23). The final genome annotation included a total of 9,627 protein-coding genes and 196 tRNAs. AntiSMASH (v4.1.0) predicted 22 putative secondary metabolite clusters (24). This annotation is comparable with that of *V. nonalfalfae* isolate T2, a lethal xylem-invading hop strain, which had 9,269 protein-coding genes and a total assembly size of 34.2 Mb (13).

Data availability. This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [RBVV00000000](https://doi.org/10.1093/nar/rbvv0000000). The version described in this paper is the first version, RBVV01000000. Sequence reads were deposited under SRA project accession number [SRP162963](https://www.ncbi.nlm.nih.gov/sra/SRP162963) and BioProject accession number [PRJNA493511](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA493511).

ACKNOWLEDGMENTS

Funding for this project was provided by USDA APHIS (agreement number AP18PPQFO000C529), USDA Forest Service, Northeastern Area, State and Private Forestry, Forest Health Technology Enterprise Team (FHTET), and United States Department of Agriculture, National Institute of Food and Agriculture hatch project CA-R-PPA-5062-H to J.E.S. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication. The genome assembly was performed at the Huck Institute, and genome annotation was performed on the High-Performance Computing Cluster at the University of California-Riverside in the Institute of Integrative Genome Biology, supported by NSF DBI-1429826 and NIH S10-OD016290.

Thanks to Richard Burhans for assistance in accessing sequence data.

REFERENCES

- Schall MJ, Davis DD. 2009. *Ailanthus altissima* wilt and mortality: etiology. *Plant Dis* 93:747–751. <https://doi.org/10.1094/PDIS-93-7-0747>.
- Kasson MT, Short DPG, O'Neal ES, Subbarao KV, Davis DD. 2014. Comparative pathogenicity, biocontrol efficacy, and multilocus sequence typing of *Verticillium nonalfalfae* from the invasive *Ailanthus altissima* and other hosts. *Phytopathology* 104:282–292. <https://doi.org/10.1094/PHYTO-06-13-0148-R>.
- Snyder AL, Salom SM, Kok LT. 2014. Survey of *Verticillium nonalfalfae* (Phyllachorales) on tree-of-heaven in the southeastern USA. *Biocontrol Sci Technol* 24:303–314. <https://doi.org/10.1080/09583157.2013.860426>.
- Rebbeck J, Malone MA, Short DPG, Kasson MT, O'Neal ES, Davis DD. 2013. First report of *Verticillium* wilt caused by *Verticillium nonalfalfae* on tree-of-heaven (*Ailanthus altissima*) in Ohio. *Plant Dis* 97:999–999. <https://doi.org/10.1094/PDIS-01-13-0062-PDN>.
- Kasson MT, Davis MD, Davis DD. 2013. The invasive *Ailanthus altissima* in Pennsylvania: a case study elucidating species introduction, migration, invasion, and growth patterns in the northeastern US. *Northeast Nat (Steuben)* 20:1–60.
- Maschek O, Halmschlager E. 2016. First report of *Verticillium* wilt on *Ailanthus altissima* in Europe caused by *Verticillium nonalfalfae*. *Plant Dis* 100:529–529. <https://doi.org/10.1094/PDIS-07-15-0733-PDN>.
- Maschek O, Halmschlager E. 2017. Natural distribution of *Verticillium* wilt on invasive *Ailanthus altissima* in eastern Austria and its potential for biocontrol. *For Pathol* 47:e12356. <https://doi.org/10.1111/efp.12356>.
- O'Neal ES, Davis DD. 2015. Biocontrol of *Ailanthus altissima*: inoculation protocol and risk assessment for *Verticillium nonalfalfae* (Plectosphaerellaceae: Phyllachorales). *Biocontrol Sci Technol* 25:950–969. <https://doi.org/10.1080/09583157.2015.1023258>.
- Schall MJ, Davis DD. 2009. *Verticillium* wilt of *Ailanthus altissima*: susceptibility of associated tree species. *Plant Dis* 93:1158–1162. <https://doi.org/10.1094/PDIS-93-11-1158>.
- Kasson MT, O'Neal ES, Davis DD. 2015. Expanded host range testing for *Verticillium nonalfalfae*: potential biocontrol agent against the invasive *Ailanthus altissima*. *Plant Dis* 99:823–835. <https://doi.org/10.1094/PDIS-04-14-0391-RE>.
- Maschek O, Halmschlager E. 2018. Effects of *Verticillium nonalfalfae* on *Ailanthus altissima* and associated indigenous and invasive tree species in eastern Austria. *Eur J Forest Res* 137:197–209. <https://doi.org/10.1007/s10342-018-1099-y>.
- O'Neal ES, Davis DD. 2015. Intraspecific root grafts and clonal growth within *Ailanthus altissima* stands influence *Verticillium nonalfalfae* transmission. *Plant Dis* 99:1070–1077. <https://doi.org/10.1094/PDIS-07-14-0722-RE>.
- Jakše J, Jelen V, Radišek S, de Jonge R, Mandelc S, Majer A, Curk T, Zupan B, Thomma BP, Javornik B. 2018. Genome sequence of a lethal strain of xylem-invading *Verticillium nonalfalfae*. *Genome Announc* 6:e01458-17. <https://doi.org/10.1128/genomeA.01458-17>.
- Song S, Kim S, Kwon SW, Lee SI, Jablonski PG. 2018. Defense sequestration associated with narrowing of diet and ontogenetic change to aposematic colours in the spotted lanternfly. *Sci Rep* 8:16831. <https://doi.org/10.1038/s41598-018-34946-y>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen Y-J, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer MLI, Jarvie TP, Jirage KB, Kim J-B, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, Lohman KL, Lu H, Makhijani VB, McDade KE, McKenna MP, Myers EW, Nickerson E, Nobile JR, Plant R, Puc BP, Ronan MT, Roth GT, Sarkis GJ, Simons JF, Simpson JW, Srinivasan M, Tartaro KR, Tomasz A, Vogt KA, Volkmer GA, Wang SH, Wang Y, Weiner MP, Yu P, Begley RF, Rothberg JM. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437:376–380. <https://doi.org/10.1038/nature03959>.
- Stajich JE, Palmer JM. 2018. AAFTE: Automated Assembly For the Fungi. <https://doi.org/10.5281/zenodo.1658103>.
- Palmer JM, Stajich JE. 2018. Funannotate: Eukaryotic Genome Annotation Pipeline. <http://funannotate.readthedocs.io>.
- Klosterman SJ, Subbarao KV, Kang S, Veronese P, Gold SE, Thomma BPHJ, Chen Z, Henrissat B, Lee Y-H, Park J, Garcia-Pedrajas MD, Barbara DJ, Anchieta A, de Jonge R, Santhanam P, Maruthachalam K, Atallah Z, Amyotte SG, Paz Z, Inderbitzin P, Hayes RJ, Heiman DI, Young S, Zeng Q, Engels R, Galagan J, Cuomo CA, Dobinson KF, Ma L-J. 2011. Comparative genomics yields insights into niche adaptation of plant vascular wilt pathogens. *PLoS Pathog* 7:e1002137. <https://doi.org/10.1371/journal.ppat.1002137>.
- Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008. Gene prediction in novel fungal genomes using an *ab initio* algorithm with unsupervised training. *Genome Res* 18:1979–1990. <https://doi.org/10.1101/gr.081612.108>.
- Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction in eukaryotes with a generalized hidden Markov model that uses hints from external sources. *BMC Bioinformatics* 7:62. <https://doi.org/10.1186/1471-2105-7-62>.
- Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31:3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>.
- Stajich JE. 2018. Fungi Gene Prediction Parameters. <https://doi.org/10.5281/zenodo.1649679>.
- Blin K, Wolf T, Chevrette MG, Lu X, Schwalen CJ, Kautsar SA, Suarez Duran HG, de Los Santos ELC, Kim HU, Nave M, Dickschat JS, Mitchell DA, Shelest E, Breitling R, Takano E, Lee SY, Weber T, Medema MH. 2017. antiSMASH 4.0—improvements in chemistry prediction and gene cluster boundary identification. *Nucleic Acids Res* 45:W36–W41. <https://doi.org/10.1093/nar/gkx319>.